



New York State
DEPARTMENT OF ENVIRONMENTAL CONSERVATION

142 2003


STR
500-6
QUACKERY
200-0183
21

Division of Water

Quackenderry Creek

Biological Assessment

2005 Survey



New York State
Department of Environmental Conservation
Region 1

George E. Pataki, Governor

Denise M. Sheehan, Commissioner

New York State Documents

OCLC:



CHECKLIST: May 2006

Original Document Scanned at:

- 400 DPI Simplex
 Duplex

Original Document contained:

- Black & White Photos
 Colored Photos
 Colored Print (list color) _____
 Colored Paper (list color) _____
 Line Art, Graphs
 Oversized Pages -- reduced from _____ (original size)
 Text Only

Date Scanned: 8/13/2008

**This electronic document has been scanned by the
New York State Library from a paper original and has been stored
on optical media.**

**The New York State Library
Cultural Education Center
Albany, NY 12230**

(MASTER.DOC. 9/99)

Quackenderry Creek
BIOLOGICAL ASSESSMENT

Lower Hudson River Basin
Rensselaer County, New York

Survey date: July 12, 2005
Report date: February 1, 2006

Robert W. Bode
Margaret A. Novak
Lawrence E. Abele
Diana L. Heitzman
Alexander J. Smith
Christine Smith


Stream Biomonitoring Unit
Bureau of Water Assessment and Management
Division of Water
NYS Department of Environmental Conservation
Albany, New York



CONTENTS

Background.....	1
Results and Conclusions.....	1
Discussion.....	2
Literature Cited.....	3
Overview of Field Data.....	3
Figure 1. Biological Assessment Profile.....	4
Table 1. Impervious Surface Calculations	4
Figure 2. Impervious Surface, Conductivity, and Nutrient Biotic Index Values	5
Table 2. Impact Source Determination.....	6
Table 3. Station Locations and Photographs	7
Figure 3. Site Location Map.....	8
Table 4. Macroinvertebrate Species Collected.....	9
Macroinvertebrate Data Reports: Raw Data	10
Field Data Summary.....	15
Laboratory Data Summary	17
Appendix I. Biological Methods for Kick Sampling.....	19
Appendix II. Macroinvertebrate Community Parameters.....	20
Appendix III. Levels of Water Quality Impact in Streams.....	21
Appendix IV. Biological Assessment Profile Description.....	22
Appendix V. Water Quality Assessment Criteria.....	24
Appendix VI. Traveling Kick Sample Illustration.....	25
Appendix VII. Macroinvertebrate Illustrations.....	26
Appendix VIII. Rationale for Biological Monitoring.....	28
Appendix IX. Glossary.....	29
Appendix X. Impact Source Determination Methods and Models.....	30
Appendix XI. Methods for Calculating the Nutrient Biotic Index	36
Appendix XII. Headwater Effects on Macroinvertebrate Communities	37
Appendix XIII. Method for Determinating of Percent Impervious Surface Cover	38





Stream: Quackenderry Creek, Rensselaer County, New York

Reach: North Greenbush to Rensselaer, New York

Drainage basin: Lower Hudson River

Background:

The Stream Biomonitoring Unit sampled Quackenderry Creek on July 21, 2005. The purpose of the sampling was to assess overall water quality, especially in relation to percent impervious surface cover of the watershed. In a riffle area at each of five sites, a traveling kick sample for macroinvertebrates was taken using methods described in the Quality Assurance document (Bode, et al., 2002) and summarized in Appendix I. The contents of each sample were field-inspected to determine major groups of organisms present, and then preserved in alcohol for laboratory inspection of a 100-specimen subsample from each site. Macroinvertebrate community parameters used in the determination of water quality included species richness, biotic index, EPT richness, and percent model affinity (see Appendices II and III). Expected variability of results is stated in Smith and Bode (2004). Table 2 provides a listing of sampling sites and Table 3 provides a listing of all macroinvertebrate species collected in the present survey. This is followed by macroinvertebrate data reports, including raw macroinvertebrate data from each site. Percent impervious surface cover was determined by Stream Biomonitoring Unit Volunteer and GIS analyst Christine Smith using methods described in Appendix XIII.

Results and Conclusions:

1. Water quality in the Quackenderry Creek ranged from non- to slightly impacted. Water quality worsened as percent impervious surface cover in the watershed increased from 15% to 21%.
2. Future increases in impervious surface cover greater than 25% are predicted to result in moderate impacts in Quackenderry Creek, and an inability to use the stream for fishing and fish propagation.

Discussion:

Quackenderry Creek originates approximately 0.7 mile east of Route 4 in North Greenbush, and flows west for 4 miles before joining the Hudson River at Rensselaer. The stream name does not appear on USGS topographic maps, but is listed in the stream gazetteer (USGS, 1981). At a former USGS gage site in Rensselaer, the drainage area is listed as 2.99 square miles. The stream is classified as C, meaning the best water use is for fishing and fish propagation. Quackenderry Creek was not previously sampled by the Stream Biomonitoring Unit.

The present sampling was conducted to document changes in macroinvertebrate communities and water quality in relation to changes in percent impervious surface cover (ISC) in a watershed. The Quackenderry Creek watershed was selected because of recent development in the basin, including the development of 34-acre parcel in 1999 into the Shoppes at Greenbush Commons shopping center. ISC was calculated for the Quackenderry Creek sites (Table 1) using techniques described in Appendix XIII.

Recent reports in the scientific literature have documented the effects of urbanization on the physical, chemical and biological characteristics of streams. A primary factor of urbanization is an increase in the percentage of ISC. Impervious surfaces are those which cover soils that previously allowed rainwater infiltration; primarily rooftops, roadways and parking areas. Effects of increasing ISC include: flooding, bank erosion, higher summer temperatures, lower winter temperatures, and increases in oxygen demand, conductivity, suspended solids, ammonium, hydrocarbons, metals, pesticides, nutrients and runoff. In examining the biological effects of elevated ISC, one proposed classification divides urban streams into three categories: sensitive (0-10% ISC), impacted (11-25% ISC), and non-supporting (26-100% ISC) (Schueler and Holland, 2000). A mitigating factor in estimating ISC is disconnected basins which provide some buffering. An example is a single-family residential area where rooftops drain to dry wells or other infiltration areas. In some calculations, these disconnected portions are subtracted from the total ISC to yield an effective ISC.

In the present study, water quality ranged from non-impacted to slightly impacted in Quackenderry Creek (Figure 1), generally declining from upstream to downstream. ISC ranged from 14% at the upstream site (Station-1) to 21% at the site immediately downstream of the shopping center (Station-3), and the ISC trend was closely correlated with specific conductance and the Nutrient Biotic Index for phosphorus (NBI-P) (Figure 2). An NBI-P value of 6.0 or greater, the provisional definition of eutrophic waters, was reported at all sites with ISC greater than 15% (Stations 2-5) in Quackenderry Creek. Impact Source Determination (Table 2) also show nutrients to be an influencing factor in the creek.

Subsequent to the building of the Shoppes at Greenbush Common, flooding occurs along Quackenderry Creek in Rensselaer downstream of Station-5 immediately after rain events, likely due to increased ISC in the basin. A dam project was developed and built one mile upstream of Station-5 to address the problem. The macroinvertebrate community at Station-5 reflects impoundment effects from the dam.

Due to the small watershed of Quackenderry Creek, two types of adjustments were made to metric values. Percent Model Affinity values at Stations 1-3 were adjusted upwards, due to high numbers of Plecoptera. Citing the Percent Model Affinity paper, "In a few cases, high contributions by an

intolerant group, usually mayflies, may result in low percent similarity values, indicating a polluted condition where one does not exist; affinity values should therefore be reviewed to determine which groups cause deviation from the model,” (Novak and Bode, 1992). The adjustment factor at each site reflects the number of Plecoptera exceeding the model.

Additionally, Stations 1-2 were adjusted for headwater effects (Appendix XII). These two sites met the criteria prescribed for headwater metric adjustment: a headwater location, a community dominated by an intolerant species, and species richness, EPT richness, or percent model affinity judged to be non-representative of actual water quality. A correction factor of 1.5 was applied to species richness and EPT richness from these two sites.

Although effects of ISC were documented in this study, Quackenderry Creek was not an ideal subject for a demonstration project. The headwater condition was a mitigating factor at upstream sites, and the most upstream site already had a high percentage of ISC, both contributing to limited fauna at the site. An ideal study situation would be a stream with a non-impacted upstream site that is not in headwater condition, and a downstream site with substantial increases in ISC.

Despite the urban/suburban setting and small size of Quackenderry Creek, many areas of the basin are still forested, and the stream maintains acceptable water quality. Future increases in ISC greater than 25% are predicted to result in moderate impacts, and an inability to use the stream for fishing and fish propagation. Many of the sites sampled in this study exhibited good habitat and high aesthetic value, yet had limited access and apparently received little use. The stream and its surrounding habitat has the potential to serve as a positive resource to the local community.

Literature Cited:

- Bode, R.W., M.A. Novak, L.E. Abele, D.L. Heitzman and A.J. Smith, 2002, Quality assurance work plan for biological stream monitoring in New York State. New York State Department of Environmental Conservation, Technical Report, 115 pages.
- Schueler, T.R, and H.K. Holland, 2000, The practice of watershed protection. Center for Watershed Protection, Ellicott City, MD. 742 pages.
- Smith, A.J. and R.W. Bode, 2004, Analysis of variability in New York State benthic macroinvertebrate samples. New York State Department of Environmental Conservation, Technical Report, 43 pages.
- U.S. Geological Survey, 1981, Drainage areas of New York streams, by river basins; A stream gazetteer; Part I - data compiled as of October 1980. U.S. Geological Survey Water-Resources Investigations, Open-file report 81-1055, 359 pages plus maps.

Overview of field data

On July 12, 2005, Quackenderry Creek at the sites sampled was 2-4 meters wide, 0.1 meters deep, and had current speeds of 40-70 cm/sec in riffles. Dissolved oxygen was 7.9-10.2 mg/l, specific conductance was 573-1642 μ mhos, pH was 7.2-7.6 and temperature was 15.7-20.8 °C (60-69 °F). Measurements for each site are found on the field data summary sheets.

Figure 1. Biological Assessment Profile of index values, Quackenderry Creek, 2005. Values are plotted on a normalized scale of water quality. The line connects the mean of the four values for each site, representing species richness, EPT richness, Hilsenhoff Biotic Index, and Percent Model Affinity. See Appendix IV for more complete explanation.




Table 1. Impervious surface calculations for Quackenderry Creek. Values reflect drainage sub-basins defined by the 5 stations.

Parameter	Station					
	1	2	3	4	5	mouth
Basin area (square meters)	891,270	1,062,257	3,128,655	4,167,136	7,127,012	7,722,131
Impervious surface area (square meters)	121,513	158,826	662,801	835,412	1,227,548	1,487,758
Percent impervious surface	13.63	14.95	21.18	20.04	17.22	19.26

Figure 2. Plot of Impervious Surface Cover, Conductivity and Nutrient Biotic Index values, Quackenderry Creek, 2005.




Table 2. Impact Source Determination, Quackenderry Creek, 2005. Numbers represent similarity to community type models for each impact category. The highest average similarities at each station are shaded. Similarities less than 50% are less conclusive. Highest numbers represent probable type of impact. See Appendix X for further explanation.

	Station				
Community Type	01	02	03	04	05
Natural: minimal human impacts	39	46	39	35	28
Nutrient enrichment	21	47	47	57	41
Toxic: industrial, municipal discharges, or urban run-off	32	32	39	58	47
Organic: sewage or animal wastes	20	31	37	44	28
Complex: municipal and/or industrial	18	31	42	39	26
Siltation	19	39	42	54	38
Impoundment	19	37	45	35	51

STATION COMMUNITY TYPE

- QUCK-01 Natural
- QUCK-02 Natural, Nutrients
- QUCK-03 Nutrients, Complex, Siltation, Impoundment
- QUCK-04 Nutrients, Toxic, Siltation
- QUCK-05 Toxic, Impoundment

Table 3. Station Locations for Quackenderry Creek,
Rensselaer County, NY, 2005


<u>STATION</u>	<u>LOCATION</u>
QUCK-01	North Greenbush, NY end of Thompson Court latitude 42°38'54" longitude 73°41'37" 3.6 river miles above mouth
QUCK-02	North Greenbush, NY off Route 43, above transmission lines latitude 42°39'04" longitude 73°42'04" 3.2 river miles above mouth
QUCK-03	North Greenbush, NY off Route 43, below runoff trib latitude 42°39'01" longitude 73°42'23" 2.9 river miles above mouth
QUCK-04	Rensselaer, NY off Ninth Street latitude 42°39'21" longitude 73°43'27" 1.7 river miles above mouth
QUCK-05	Rensselaer, NY Below Wilson Street bridge latitude 42°38'40" longitude 73°44'09" 0.6 river miles above mouth



Figure 3



Site Overview Map

Quackenderry Creek



Legend

- Hydrography
- Drainage Boundary
- Impervious Surface



Water Quality Assessment based on Resident Macroinvertebrates

- non-impacted
- ▲ slightly impacted
- moderately impacted
- ◆ severely impacted

Table 4. Macroinvertebrates Species collected in Quackenderry Creek, July 12, 2005

PLATYHELMINTHES	
TURBELLARIA	
Planariidae	
Undetermined Turbellaria	
OLIGOCHAETA	
LUMBRICIDA	
Undetermined Lumbricina	
LUMBRICULIDA	
Lumbriculidae	
Undetermined Lumbriculidae	
TUBIFICIDA	
Enchytraeidae	
Undetermined Enchytraeidae	
Tubificidae	
Undet. Tubificidae w/ cap. setae	
Undet. Tubificidae w/o cap. setae	
MOLLUSCA	
PELECYPODA	
Sphaeriidae	
<i>Pisidium</i> sp.	
ARTHROPODA	
CRUSTACEA	
AMPHIPODA	
Gammaridae	
<i>Gammarus</i> sp.	
ISOPODA	
Asellidae	
<i>Caecidotea racovitzai</i>	
INSECTA	
EPHEMEROPTERA	
Baetidae	
<i>Baetis flavistriga</i>	
PLECOPTERA	
Leuctridae	
<i>Leuctra</i> sp.	
ODONATA	
Aeschnidae	
<i>Boyeria</i> sp.	
COLEOPTERA	
Psephenidae	
<i>Ectopria nervosa</i>	
<i>Psephenus herricki</i>	
Elmidae	
<i>Macronychus glabratus</i>	
<i>Optioservus fastiditus</i>	
<i>Stenelmis crenata</i>	
<i>Stenelmis</i> sp.	
MEGALOPTERA	
Sialidae	
<i>Sialis</i> sp.	
TRICHOPTERA	
Philopotamidae	
<i>Chimarra aterrima?</i>	
<i>Dolophilodes</i> sp.	
	Hydropsychidae
	<i>Cheumatopsyche</i> sp.
	<i>Hydropsyche betteni</i>
	<i>Hydropsyche slossonae</i>
	<i>Hydropsyche sparna</i>
	<i>Potamyia</i> sp.
	Rhyacophilidae
	<i>Rhyacophila</i> sp.
	DIPTERA
	Tipulidae
	<i>Antocha</i> sp.
	<i>Dicranota</i> sp.
	<i>Hexatoma</i> sp.
	<i>Tipula</i> sp.
	Simuliidae
	<i>Simulium tuberosum</i>
	<i>Simulium vittatum</i>
	<i>Simulium</i> sp.
	Athericidae
	<i>Atherix</i> sp.
	Empididae
	<i>Hemerodromia</i> sp.
	Chironomidae
	<i>Natarsia</i> sp. A
	<i>Thienemannimyia</i> gr. spp.
	<i>Diamesa</i> sp.
	<i>Pagastia orthogonia</i>
	<i>Brillia</i> sp.
	<i>Cricotopus bicinctus</i>
	<i>Cricotopus tremulus</i> gr.
	<i>Eukiefferiella claripennis</i> gr.
	<i>Parametrioctenemus lundbecki</i>
	<i>Tvetenia bavarica</i> gr.
	<i>Polypedilum aviceps</i>
	<i>Polypedilum illinoense</i>
	<i>Polypedilum tuberculum</i>
	<i>Paratanytarsus</i> sp.
	<i>Rheotanytarsus exiguus</i> gr.

Macroinvertebrate Data Reports: Raw Data

STREAM SITE: Quackenderry Creek, Station QUICK- 01
 LOCATION: North Greenbush, NY, off Thompson Court
 DATE: 12 July 2005
 SAMPLE TYPE: Kick sample
 SUBSAMPLE: 100 organisms

ANNELIDA

OLIGOCHAETA

TUBIFICIDA	Tubificidae	Undet. Tubificidae w/o cap. setae	1
------------	-------------	-----------------------------------	---

ARTHROPODA

INSECTA

PLECOPTERA	Leuctridae	<i>Leuctra sp.</i>	29
------------	------------	--------------------	----

COLEOPTERA	Psephenidae	<i>Ectopria nervosa</i>	1
------------	-------------	-------------------------	---

TRICHOPTERA	Philopotamidae	<i>Chimarra aterrima?</i>	1
-------------	----------------	---------------------------	---

	Hydropsychidae	<i>Potamyia sp.</i>	8
--	----------------	---------------------	---

DIPTERA	Tipulidae	<i>Dicranota sp.</i>	35
---------	-----------	----------------------	----

		<i>Tipula sp.</i>	2
--	--	-------------------	---

	Empididae	<i>Hemerodromia sp.</i>	1
--	-----------	-------------------------	---

	Chironomidae	<i>Thienemannimyia gr. spp.</i>	1
--	--------------	---------------------------------	---

		<i>Pagastia orthogonia</i>	1
--	--	----------------------------	---

		<i>Tvetenia bavarica gr.</i>	2
--	--	------------------------------	---

		<i>Polypedilum aviceps</i>	9
--	--	----------------------------	---

		<i>Polypedilum tuberculum</i>	9
--	--	-------------------------------	---

SPECIES RICHNESS: 13 (good*)
 BIOTIC INDEX: 2.87 (very good)
 EPT RICHNESS: 3 (poor*)
 MODEL AFFINITY: 46 (very good**)
 NUTRIENT INDEX (P) 5.44 (oligotrophic)
 ASSESSMENT: slightly impacted (6.88)

DESCRIPTION: This site was approximately 0.4 mile from the stream source. The habitat was well-shaded, but the stream was silty and slow-moving. The macroinvertebrate community was heavily dominated by crane-fly larvae and stoneflies. The indication of slight impact was likely due to the slow-moving nature of the stream. Nearly all the species present were considered intolerant.

* Metrics were adjusted due to headwater conditions. See Appendix XII.

** Percent Model Affinity was adjusted up by the percent contribution of Plecoptera exceeding the model.

Macroinvertebrate Data Reports: Raw Data (cont.)

STREAM SITE: Quackenderry Creek, Station QUICK- 02
 LOCATION: North Greenbush, NY, off Route 43
 DATE: 12 July 2005
 SAMPLE TYPE: Kick sample
 SUBSAMPLE: 100 organisms

ANNELIDA

OLIGOCHAETA

TUBIFICIDA	Tubificidae	Undet. Tubificidae w/o cap. setae	1
------------	-------------	-----------------------------------	---

ARTHROPODA

INSECTA

EPHEMEROPTERA	Baetidae	<i>Baetis intercalaris</i>	1
PLECOPTERA	Leuctridae	<i>Leuctra sp.</i>	30
COLEOPTERA	Psephenidae	<i>Psephenus herricki</i>	1
		<i>Elmidae</i>	16
		<i>Stenelmis crenata</i>	4
TRICHOPTERA	Philopotamidae	<i>Dolophilodes sp.</i>	1
		<i>Hydropsyche sparna</i>	1
		<i>Potamyia sp.</i>	25
DIPTERA	Tipulidae	<i>Antocha sp.</i>	1
		<i>Dicranota sp.</i>	8
		<i>Tipula sp.</i>	3
	Chironomidae	<i>Thienemannimyia gr. spp.</i>	1
		<i>Pagastia orthogonia</i>	2
		<i>Tvetenia bavarica gr.</i>	1
		<i>Polypedilum aviceps</i>	4

SPECIES RICHNESS: 16 (good*)
 BIOTIC INDEX: 3.07 (very good)
 EPT RICHNESS: 8 (good*)
 MODEL AFFINITY: 45 (very good**)
 NUTRIENT INDEX (P) 6.22 (eutrophic)
 ASSESSMENT: non-impacted (7.53)

DESCRIPTION: The sampling site was off Route 43 in North Greenbush, upstream of overhead transmission lines. Habitat was good, with faster current than Station-1. Based on adjusted values, water quality was assessed as non-impacted. The NBI-P indicated increased nutrients compared to Station-1.

* Metrics were adjusted due to headwater conditions. See Appendix XII.

** Percent Model Affinity was adjusted up by the percent contribution of Plecoptera exceeding the model.

Macroinvertebrate Data Reports: Raw Data (cont.)

STREAM SITE: Quackenderry Creek, Station QUICK- 03
 LOCATION: North Greenbush, NY, off Route 43
 DATE: 12 July 2005
 SAMPLE TYPE: Kick sample
 SUBSAMPLE: 100 organisms

PLATYHELMINTHES

TURBELLARIA

Planariidae	Undetermined Turbellaria	1
-------------	--------------------------	---

ANNELIDA

OLIGOCHAETA

TUBIFICIDA

Enchytraeidae	Undetermined Enchytraeidae	1
Tubificidae	Undet. Tubificidae w/ cap. setae	1
	Undet. Tubificidae w/o cap. setae	11

ARTHROPODA

INSECTA

EPHEMEROPTERA

PLECOPTERA

COLEOPTERA

MEGALOPTERA

TRICHOPTERA

DIPTERA

Baetidae	<i>Baetis intercalaris</i>	4
Leuctridae	<i>Leuctra sp.</i>	9
Elmidae	<i>Optioservus fastiditus</i>	2
	<i>Stenelmis sp.</i>	7
Sialidae	<i>Sialis sp.</i>	1
Hydropsychidae	<i>Hydropsyche betteni</i>	5
	<i>Potamyia sp.</i>	20
Rhyacophilidae	<i>Rhyacophila sp.</i>	1
Tipulidae	<i>Dicranota sp.</i>	3
	<i>Hexatoma sp.</i>	1
Simuliidae	<i>Simulium tuberosum</i>	1
Chironomidae	<i>Diamesa sp.</i>	20
	<i>Pagastia orthogonia</i>	4
	<i>Prodiamesa olivacea</i>	1
	<i>Brillia sp.</i>	1
	<i>Cricotopus bicinctus</i>	1
	<i>Tvetenia bavarica gr.</i>	2
	<i>Polypedilum aviceps</i>	2
	<i>Paratanytarsus sp.</i>	1

SPECIES RICHNESS: 23 (good)
 BIOTIC INDEX: 5.00 (good)
 EPT RICHNESS: 5 (poor)
 MODEL AFFINITY: 60 (good*)
 NUTRIENT INDEX (P): 7.26 (eutrophic)
 ASSESSMENT: slightly impacted (6.37)

DESCRIPTION: This site was only 0.25 mile downstream of Station-2, but received much more drainage, including run-off from the Shoppes at Greenbush Common. Habitat was comparable to that at Station-2. Conductivity had increased from 683 to 1642 μ mhos, and the water appeared grey. The macroinvertebrate fauna had changed substantially from Station-2, being dominated by facultative midges and caddisflies; stoneflies were much less numerous. Using adjusted metric values, all values worsened compared to Station-2. Overall water quality was assessed as slightly impacted.

* Percent Model Affinity was adjusted up by the percent contribution of Plecoptera exceeding the model.

Macroinvertebrate Data Reports: Raw Data (cont.)

STREAM SITE: Quackenderry Creek, Station QUICK- 04
 LOCATION: Rensselaer, NY, off Ninth Street
 DATE: 12 July 2005
 SAMPLE TYPE: Kick sample
 SUBSAMPLE: 100 organisms

PLATYHELMINTHES

TURBELLARIA

Planariidae	Undetermined Turbellaria	1
-------------	--------------------------	---

ARTHROPODA

CRUSTACEA

ISOPODA	Asellidae	<i>Caecidotea racovitzai</i>	1
---------	-----------	------------------------------	---

INSECTA

EPHEMEROPTERA	Baetidae	<i>Baetis flavistriga</i>	1
---------------	----------	---------------------------	---

COLEOPTERA	Elmidae	<i>Optioservus fastiditus</i>	2
------------	---------	-------------------------------	---

		<i>Stenelmis crenata</i>	24
--	--	--------------------------	----

TRICHOPTERA	Philopotamidae	<i>Chimarra aterrima?</i>	5
-------------	----------------	---------------------------	---

	Hydropsychidae	<i>Cheumatopsyche sp.</i>	8
--	----------------	---------------------------	---

		<i>Hydropsyche betteni</i>	4
--	--	----------------------------	---

		<i>Hydropsyche slossonae</i>	12
--	--	------------------------------	----

		<i>Hydropsyche sparna</i>	3
--	--	---------------------------	---

DIPTERA	Tipulidae	<i>Dicranota sp.</i>	23
---------	-----------	----------------------	----

	Simuliidae	<i>Simulium tuberosum</i>	1
--	------------	---------------------------	---

		<i>Atherix sp.</i>	2
--	--	--------------------	---

	Chironomidae	<i>Natarsia sp. A</i>	1
--	--------------	-----------------------	---

		<i>Thienemannimyia gr. spp.</i>	3
--	--	---------------------------------	---

		<i>Pagastia orthogonia</i>	1
--	--	----------------------------	---

		<i>Cricotopus bicinctus</i>	1
--	--	-----------------------------	---

		<i>Cricotopus tremulus gr.</i>	4
--	--	--------------------------------	---

		<i>Parametriocnemus lundbecki</i>	1
--	--	-----------------------------------	---

		<i>Polypedilum tuberculum</i>	1
--	--	-------------------------------	---

		<i>Rheotanytarsus exiguus gr.</i>	1
--	--	-----------------------------------	---

SPECIES RICHNESS: 21 (good)
 BIOTIC INDEX: 4.60 (good)
 EPT RICHNESS: 6 (good)
 MODEL AFFINITY: 44 (poor)
 NUTRIENT INDEX (P): 6.02 (eutrophic)
 ASSESSMENT: slightly impacted (5.71)

DESCRIPTION: The sampling site was accessed down a steep slope at Ninth Street and Birch Street in Rensselaer. Habitat was considered acceptable, and comparable to upstream sites. The macroinvertebrate community was dominated by caddisflies, riffle beetles, and crane fly larvae. Based on the metrics, water quality was assessed as slightly impacted.

Macroinvertebrate Data Reports: Raw Data (cont.)

STREAM SITE: Quackenderry Creek, Station QUICK- 05
 LOCATION: Rensselaer, NY, below Wilson Street bridge
 DATE: 12 July 2005
 SAMPLE TYPE: Kick sample
 SUBSAMPLE: 100 organisms

ANNELIDA

OLIGOCHAETA

TUBIFICIDA	Tubificidae	Undet. Tubificidae w/o cap. setae	8
------------	-------------	-----------------------------------	---

ARTHROPODA

CRUSTACEA

AMPHIPODA	Gammaridae	<i>Gammarus sp.</i>	1
-----------	------------	---------------------	---

INSECTA

EPHEMEROPTERA	Baetidae	<i>Baetis flavistriga</i>	2
---------------	----------	---------------------------	---

ODONATA	Aeschnidae	<i>Boyeria sp.</i>	2
---------	------------	--------------------	---

COLEOPTERA	Psephenidae	<i>Psephenus herricki</i>	1
------------	-------------	---------------------------	---

Elmidae	<i>Macronychus glabratus</i>	3
---------	------------------------------	---

	<i>Stenelmis crenata</i>	30
--	--------------------------	----

MEGALOPTERA	Sialidae	<i>Sialis sp.</i>	1
-------------	----------	-------------------	---

TRICHOPTERA	Hydropsychidae	<i>Cheumatopsyche sp.</i>	4
-------------	----------------	---------------------------	---

	<i>Hydropsyche slossonae</i>	2
--	------------------------------	---

DIPTERA	Tipulidae	<i>Antocha sp.</i>	1
---------	-----------	--------------------	---

	<i>Dicranota sp.</i>	11
--	----------------------	----

Simuliidae	<i>Simulium vittatum</i>	2
------------	--------------------------	---

	<i>Simulium sp.</i>	1
--	---------------------	---

Athericidae	<i>Atherix sp.</i>	6
-------------	--------------------	---

Chironomidae	<i>Natarsia sp. A</i>	2
--------------	-----------------------	---

	<i>Thienemannimyia gr. spp.</i>	8
--	---------------------------------	---

	<i>Diamesa sp.</i>	3
--	--------------------	---

	<i>Pagastia orthogonia</i>	4
--	----------------------------	---

	<i>Eukiefferiella claripennis gr.</i>	2
--	---------------------------------------	---

	<i>Parametriocnemus lundbecki</i>	4
--	-----------------------------------	---

	<i>Polypedilum illinoense</i>	2
--	-------------------------------	---

SPECIES RICHNESS: 22 (good)
 BIOTIC INDEX: 5.11 (good)
 EPT RICHNESS: 3 (poor)
 MODEL AFFINITY: 53 (good)
 NUTRIENT INDEX (P): 6.95 (eutrophic)
 ASSESSMENT: slightly impacted (5.55)

DESCRIPTION: The kick sample was taken 20 meters downstream of the Wilson Street bridge in Rensselaer. It was 0.95 mile downstream of a dam on Quackenderry Creek. The site had much urban refuse in the stream, and abundant brown algae and silt were also present. The macroinvertebrate community was dominated by facultative riffle beetles and midges, and water quality was assessed as slightly impacted. Impact Source Determination reflected impoundment and urban runoff.

FIELD DATA SUMMARY

STREAM NAME: Quackenderry Creek		DATE SAMPLED: 7/12/2005		
REACH: North Greenbush to Rensselaer				
FIELD PERSONNEL INVOLVED: Bode, Smith				
STATION	01	02	03	04
ARRIVAL TIME AT STATION	8:00 AM	8:40 AM	9:10 AM	9:45 AM
LOCATION	North Greenbush off Thompson Ct	North Greenbush off Exit 8 ramp	Rensselaer off Exit 8 ramp	Rensselaer Ninth & Birch St
PHYSICAL CHARACTERISTICS				
Width (meters)	2.0	2.0	3.0	3.0
Depth (meters)	0.1	0.1	0.1	0.1
Current speed (cm per sec.)	40	70	70	70
Substrate (%)				
Rock (>25.4 cm, or bedrock)	10	10	20	10
Rubble (6.35 – 25.4 cm)	40	40	30	30
Gravel (0.2 – 6.35 cm)	20	20	20	30
Sand (0.06 – 2.0 mm)	10	10	10	10
Silt (0.004 – 0.06 mm)	20	20	20	20
Embeddedness (%)	30	30	20	30
CHEMICAL MEASUREMENTS				
Temperature (° C)	19.0	19.5	16.7	20.8
Specific Conductance (umhos)	573	683	1642	1609
Dissolved Oxygen (mg/l)	7.9	8.4	10.2	8.1
pH	7.2	7.2	7.5	7.6
BIOLOGICAL ATTRIBUTES				
Canopy (%)	100	50	70	90
Aquatic Vegetation				
algae – suspended				
algae – attached, filamentous				
algae – diatoms				
macrophytes or moss				
Occurrence of Macroinvertebrates				
Ephemeroptera (mayflies)			x	x
Plecoptera (stoneflies)	x	x	x	
Trichoptera (caddisflies)	x	x	x	x
Coleoptera (beetles)				x
Megaloptera (dobsonflies, alderflies)				x
Odonata (dragonflies, damselflies)		x		x
Chironomidae (midges)	x	x	x	x
Simuliidae (black flies)				
Decapoda (crayfish)	x	x	x	
Gammaridae (scuds)				
Mollusca (snails, clams)				
Oligochaeta (worms)	x	x		
Other		x		x
FAUNAL CONDITION	Very good	Very good	Very good	Good

FIELD DATA SUMMARY

STREAM NAME: Quackenderry Creek **DATE SAMPLED:** 7/12/2005

REACH: North Greenbush to Rensselaer

FIELD PERSONNEL INVOLVED: Bode, Smith

STATION	05			
ARRIVAL TIME AT STATION	10:20 AM			
LOCATION	Rensselaer Wilson St bridge			
PHYSICAL CHARACTERISTICS				
Width (meters)	4.0			
Depth (meters)	0.1			
Current speed (cm per sec.)	70			
Substrate (%)				
Rock (>25.4 cm, or bedrock)	10			
Rubble (6.35 – 25.4 cm)	30			
Gravel (0.2 – 6.35 cm)	30			
Sand (0.06 – 2.0 mm)	10			
Silt (0.004 – 0.06 mm)	20			
Embeddedness (%)	30			
CHEMICAL MEASUREMENTS				
Temperature (° C)	15.7			
Specific Conductance (umhos)	1107			
Dissolved Oxygen (mg/l)	9.4			
pH	7.5			
BIOLOGICAL ATTRIBUTES				
Canopy (%)	80			
Aquatic Vegetation				
algae – suspended				
algae – attached, filamentous				
algae – diatoms				
macrophytes or moss				
Occurrence of Macroinvertebrates				
Ephemeroptera (mayflies)	x			
Plecoptera (stoneflies)				
Trichoptera (caddisflies)	x			
Coleoptera (beetles)	x			
Megaloptera (dobsonflies, alderflies)				
Odonata (dragonflies, damselflies)	x			
Chironomidae (midges)	x			
Simuliidae (black flies)				
Decapoda (crayfish)				
Gammaridae (scuds)				
Mollusca (snails, clams)				
Oligochaeta (worms)				
Other	x			
FAUNAL CONDITION	Good			

LABORATORY DATA SUMMARY

STREAM NAME: Quackenderry Creek		DRAINAGE: 13			
DATE SAMPLED: 7/12/2005		COUNTY: Rensselaer			
SAMPLING METHOD: Travelling Kick					
STATION LOCATION	01 North Greenbush off Thompson Ct	02 North Greenbush off Exit 8 ramp	03 Rensselaer off Exit 8 ramp	04 Rensselaer Ninth & Birch St	
DOMINANT SPECIES/% CONTRIBUTION/TOLERANCE/COMMON NAME					
	1.	Dicronata sp. 35 % intolerant crane fly	Leuctra sp. 30 % intolerant stone fly	Potamyia sp. 20 % intolerant caddisfly	Stenelmis crenata 24 % facultative beetle
Intolerant = not tolerant of poor water quality	2.	Leuctra sp. 29 % intolerant stone fly	Potamyia sp. 25 % intolerant caddisfly	Diamesa sp. 20 % facultative midge	Dicronata sp. 23 % intolerant crane fly
Facultative = occurring over a wide range of water quality	3.	Polypedilum aviceps 9 % facultative midge	Optioservus fastiditus 16% intolerant beetle	Undet. Tubificidae w/o cap. setae 11 % facultative worm	Hydropsyche slossonae 12 % facultative caddisfly
Tolerant = tolerant of poor water quality	4.	Polypedilum tuberculum 9 % facultative midge	Dicronata sp. 8 % intolerant crane fly	Leuctra sp. 9 % intolerant stone fly	Cheumatopsyche sp. 8 % facultative caddisfly
	5.	Potamyia sp. 8 % intolerant caddisfly	Stenelmis crenata 4 % facultative beetle	Stenelmis sp. 7 % facultative beetle	Chimarra aterrima? 5 % intolerant caddisfly
% CONTRIBUTION OF MAJOR GROUPS (NUMBER OF TAXA IN PARENTHESES)					
Chironomidae (midges)		22.0 (5)	8.0 (4)	32.0 (8)	13.0 (8)
Trichoptera (caddisflies)		9.0 (2)	27.0 (3)	26.0 (3)	32.0 (5)
Ephemeroptera (mayflies)		0.0 (0)	1.0 (1)	4.0 (1)	1.0 (1)
Plecoptera (stoneflies)		29.0 (1)	30.0 (1)	9.0 (1)	0.0 (0)
Coleoptera (beetles)		1.0 (1)	21.0 (3)	9.0 (2)	26.0 (2)
Oligochaeta (worms)		1.0 (1)	1.0 (1)	13.0 (3)	0.0 (0)
Mollusca (clams and snails)		0.0 (0)	0.0 (0)	0.0 (0)	0.0 (0)
Crustacea (crayfish, scuds, sowbugs)		0.0 (0)	0.0 (0)	0.0 (0)	1.0 (1)
Other insects (odonates, diptera)		38.0 (3)	12.0 (3)	6.0 (4)	26.0 (3)
Other (Nemertea, Platyhelminthes)		0.0 (0)	0.0 (0)	1.0 (1)	1.0 (1)
SPECIES RICHNESS		20	24	23	21
BIOTIC INDEX		2.87	3.07	5.00	4.60
EPT RICHNESS		5	8	5	6
PERCENT MODEL AFFINITY		70	70	64	44
FIELD ASSESSMENT		Very good	Very good	Very good	Good
OVERALL ASSESSMENT		Slightly impacted	Non-impacted	Slightly impacted	Slightly impacted

LABORATORY DATA SUMMARY

STREAM NAME: Quackenderry Creek		DRAINAGE: 13		
DATE SAMPLED: 7/12/2005		COUNTY: Rensselaer		
SAMPLING METHOD: Travelling Kick				
STATION LOCATION	05 Rensselaer Wilson St bridge			
DOMINANT SPECIES/% CONTRIBUTION/TOLERANCE/COMMON NAME				
	1.	Stenelmis crenata 30 % facultative beetle		
Intolerant = not tolerant of poor water quality	2.	Dicronata sp. 11 % intolerant crane fly		
Facultative = occurring over a wide range of water quality	3.	Undet. Tubificidae w/o cap. setae 8 % facultative worm		
Tolerant = tolerant of poor water quality	4.	Thienemannimyia gr. spp. 8 % facultative midge		
	5.	Atherix sp. 6% intolerant crane fly		
% CONTRIBUTION OF MAJOR GROUPS (NUMBER OF TAXA IN PARENTHESES)				
Chironomidae (midges)		25.0 (7)		
Trichoptera (caddisflies)		6.0 (2)		
Ephemeroptera (mayflies)		2.0 (1)		
Plecoptera (stoneflies)		0.0 (0)		
Coleoptera (beetles)		34.0 (3)		
Oligochaeta (worms)		8.0 (1)		
Mollusca (clams and snails)		0.0 (0)		
Crustacea (crayfish, scuds, sowbugs)		1.0 (1)		
Other insects (odonates, diptera)		24.0 (7)		
Other (Nemertea, Platyhelminthes)		0.0 (0)		
SPECIES RICHNESS		22		
BIOTIC INDEX		5.11		
EPT RICHNESS		3		
PERCENT MODEL AFFINITY		53		
FIELD ASSESSMENT		Good		
OVERALL ASSESSMENT		Slightly impacted		

Appendix I. Biological Methods for Kick Sampling

A. Rationale. The use of the standardized kick sampling method provides a biological assessment technique that lends itself to rapid assessments of stream water quality.

B. Site Selection. Sampling sites are selected based on these criteria: (1) The sampling location should be a riffle with a substrate of rubble, gravel and sand; depth should be one meter or less, and current speed should be at least 0.4 meters per second. (2) The site should have comparable current speed, substrate type, embeddedness, and canopy cover to both upstream and downstream sites to the degree possible. (3) Sites are chosen to have a safe and convenient access.

C. Sampling. Macroinvertebrates are sampled using the standardized traveling kick method. An aquatic net is positioned in the water at arms' length downstream and the stream bottom is disturbed by foot, so that organisms are dislodged and carried into the net. Sampling is continued for a specified time and distance in the stream. Rapid assessment sampling specifies sampling for five minutes over a distance of five meters. The contents of the net are emptied into a pan of stream water. The contents are then examined, and the major groups of organisms are recorded, usually on the ordinal level (e.g., stoneflies, mayflies, caddisflies). Larger rocks, sticks, and plants may be removed from the sample if organisms are first removed from them. The contents of the pan are poured into a U.S. No. 30 sieve and transferred to a quart jar. The sample is then preserved by adding 95% ethyl alcohol.

D. Sample Sorting and Subsampling. In the laboratory, the sample is rinsed with tap water in a U.S. No. 40 standard sieve to remove any fine particles left in the residues from field sieving. The sample is transferred to an enamel pan and distributed homogeneously over the bottom of the pan. A small amount of the sample is randomly removed with a spatula, rinsed with water, and placed in a petri dish. This portion is examined under a dissecting stereomicroscope and 100 organisms are randomly removed from the debris. As they are removed, they are sorted into major groups, placed in vials containing 70 percent alcohol, and counted. The total number of organisms in the sample is estimated by weighing the residue from the picked subsample and determining its proportion of the total sample weight.

E. Organism Identification. All organisms are identified to the species level whenever possible. Chironomids and oligochaetes are slide-mounted and viewed through a compound microscope; most other organisms are identified as whole specimens using a dissecting stereomicroscope. The number of individuals in each species, and the total number of individuals in the subsample is recorded on a data sheet. All organisms from the subsample are archived (either slide-mounted or preserved in alcohol). If the results of the identification process are ambiguous, suspected of being spurious, or do not yield a clear water quality assessment, additional subsampling may be required.

Appendix II. Macroinvertebrate Community Parameters

1. Species Richness is the total number of species or taxa found in the sample. For subsamples of 100-organisms each that are taken from kick samples, expected ranges in most New York State streams are: greater than 26, non-impacted; 19-26, slightly impacted; 11-18, moderately impacted; less than 11, severely impacted.

2. EPT Richness denotes the total number of species of mayflies (Ephemeroptera), stoneflies (Plecoptera), and caddisflies (Trichoptera) found in an average 100-organisms subsample. These are considered to be clean-water organisms, and their presence is generally correlated with good water quality (Lenat, 1987). Expected assessment ranges from most New York State streams are: greater than 10, non-impacted; 6-10, slightly impacted; 2-5, moderately impacted; and 0-1, severely impacted.

3. Hilsenhoff Biotic Index is a measure of the tolerance of organisms in a sample to organic pollution (sewage effluent, animal wastes) and low dissolved oxygen levels. It is calculated by multiplying the number of individuals of each species by its assigned tolerance value, summing these products, and dividing by the total number of individuals. On a 0-10 scale, tolerance values range from intolerant (0) to tolerant (10). For the purpose of characterizing species' tolerance, intolerant = 0-4, facultative = 5-7, and tolerant = 8-10. Tolerance values are listed in Hilsenhoff (1987). Additional values are assigned by the NYS Stream Biomonitoring Unit. The most recent values for each species are listed in Quality Assurance document, Bode et al. (2002). Impact ranges are: 0-4.50, non-impacted; 4.51-6.50, slightly impacted; 6.51-8.50, moderately impacted; and 8.51-10.00, severely impacted.

4. Percent Model Affinity is a measure of similarity to a model, non-impacted community based on percent abundance in seven major macroinvertebrate groups (Novak and Bode, 1992). Percent abundances in the model community are: 40% Ephemeroptera; 5% Plecoptera; 10% Trichoptera; 10% Coleoptera; 20% Chironomidae; 5% Oligochaeta; and 10% Other. Impact ranges are: greater than 64, non-impacted; 50-64, slightly impacted; 35-49, moderately impacted; and less than 35, severely impacted.

Bode, R.W., M.A. Novak, L.E. Abele, D. L. Heitzman and A.J. Smith, 2002,. Quality assurance work plan for biological stream monitoring in New York State. NYSDEC Technical Report, 115 pages.

Hilsenhoff, W. L., 1987, An improved biotic index of organic stream pollution. *The Great Lakes Entomologist* 20(1): 31-39.

Lenat, D. R., 1987, Water quality assessment using a new qualitative collection method for freshwater benthic macroinvertebrates. North Carolina Division of Environmental Management Technical Report, 12 pages.

Novak, M.A., and R.W. Bode, 1992, Percent model affinity: a new measure of macroinvertebrate community composition. *J. N. Am. Benthol. Soc.* 11(1): 80-85.

Appendix III. Levels of Water Quality Impact in Streams

The description of overall stream water quality based on biological parameters uses a four-tiered system of classification. Level of impact is assessed for each individual parameter and then combined for all parameters to form a consensus determination. Four parameters are used: species richness, EPT richness, biotic index, and percent model affinity (see Appendix II). The consensus is based on the determination of the majority of the parameters. Since parameters measure different aspects of the macroinvertebrate community, they cannot be expected to always form unanimous assessments. The assessment ranges given for each parameter are based on subsamples of 100-organisms each that are taken from macroinvertebrate riffle kick samples. These assessments also apply to most multiplate samples, with the exception of percent model affinity.

1. *Non-impacted* Indices reflect very good water quality. The macroinvertebrate community is diverse, usually with at least 27 species in riffle habitats. Mayflies, stoneflies, and caddisflies are well-represented; the EPT richness is greater than 10. The biotic index value is 4.50 or less. Percent model affinity is greater than 64. Water quality should not be limiting to fish survival or propagation. This level of water quality includes both pristine habitats and those receiving discharges which minimally alter the biota.


2. *Slightly impacted* Indices reflect good water quality. The macroinvertebrate community is slightly but significantly altered from the pristine state. Species richness usually is 19-26. Mayflies and stoneflies may be restricted, with EPT richness values of 6-10. The biotic index value is 4.51-6.50. Percent model affinity is 50-64. Water quality is usually not limiting to fish survival, but may be limiting to fish propagation.

3. *Moderately impacted* Indices reflect poor water quality. The macroinvertebrate community is altered to a large degree from the pristine state. Species richness usually is 11-18 species. Mayflies and stoneflies are rare or absent, and caddisflies are often restricted; the EPT richness is 2-5. The biotic index value is 6.51-8.50. The percent model affinity value is 35-49. Water quality often is limiting to fish propagation, but usually not to fish survival.

4. *Severely impacted* Indices reflect very poor water quality. The macroinvertebrate community is limited to a few tolerant species. Species richness is 10 or less. Mayflies, stoneflies, and caddisflies are rare or absent; EPT richness is 0-1. The biotic index value is greater than 8.50. Percent model affinity is less than 35. The dominant species are almost all tolerant, and are usually midges and worms. Often 1-2 species are very abundant. Water quality is often limiting to both fish propagation and fish survival.

Appendix IV-A. Biological Assessment Profile: Conversion of Index Values to Common 10-Scale

The Biological Assessment Profile of index values, developed by Phil O'Brien, Division of Water, NYSDEC, is a method of plotting biological index values on a common scale of water quality impact. Values from the four indices defined in Appendix II are converted to a common 0-10 scale using the formulae in the Quality Assurance document (Bode et al.,2002), and as shown in the figure below.



Appendix IV-B. Biological Assessment Profile: Plotting Values


To plot survey data:

1. Position each site on the x-axis according to miles or tenths of a mile upstream of the mouth.
2. Plot the values of the four indices for each site as indicated by the common scale.
3. Calculate the mean of the four values and plot the result. This represents the assessed impact for each site.

Example data:

	Station 1		Station 2	
	metric value	10-scale value	metric value	10-scale value
Species richness	20	5.59	33	9.44
Hilsenhoff biotic index	5.00	7.40	4.00	8.00
EPT richness	9	6.80	13	9.00
Percent model affinity	55	5.97	65	7.60
Average		6.44 (slight)		8.51 (non-)

Table IV-B. Sample Plot of Biological Assessment Profile values



Appendix V. Water Quality Assessment Criteria

Water Quality Assessment Criteria for Non-Navigable Flowing Waters

	Species Richness	Hilsenhoff Biotic Index	EPT Richness	Percent Model Affinity#	Species Diversity*
Non-Impacted	>26	0.00-4.50	>10	>64	>4
Slightly Impacted	19-26	4.51-6.50	6-10	50-64	3.01-4.00
Moderately Impacted	11-18	6.51-8.50	2-5	35-49	2.01-3.00
Severely Impacted	0-10	8.51-10.00	0-1	<35	0.00-2.00

Percent model affinity criteria are used for traveling kick samples but not for multiplate samples.


* Diversity criteria are used for multiplate samples but not for traveling kick samples.

Water Quality Assessment Criteria for Navigable Flowing Waters

	Species Richness	Hilsenhoff Biotic Index	EPT Richness	Species Diversity
Non-Impacted	>21	0.00-7.00	>5	>3.00
Slightly Impacted	17-21	7.01-8.00	4-5	2.51-3.00
Moderately Impacted	12-16	8.01-9.00	2-3	2.01-2.50
Severely Impacted	0-11	9.01-10.00	0-1	0.00-2.00

Appendix VI.


The Traveling Kick Sample



Rocks and sediment in a riffle are dislodged by foot upstream of a net. Dislodged organisms are carried by the current into the net. Sampling continues for five minutes, as the sampler gradually moves downstream to cover a distance of five meters.


Appendix VII. A.
Aquatic Macroinvertebrates that Usually Indicate Good Water Quality

Mayfly nymphs are often the most numerous organisms found in clean streams. They are sensitive to most types of pollution, including low dissolved oxygen (less than 5 ppm), chlorine, ammonia, metals, pesticides, and acidity. Most mayflies are found clinging to the undersides of rocks.




MAYFLIES

Stonefly nymphs are mostly limited to cool, well-oxygenated streams. They are sensitive to most of the same pollutants as mayflies, except acidity. They are usually much less numerous than mayflies. The presence of even a few stoneflies in a stream suggests that good water quality has been maintained for several months.




STONEFLIES

Caddisfly larvae often build a portable case of sand, stones, sticks, or other debris. Many caddisfly larvae are sensitive to pollution, although a few are tolerant. One family spins nets to catch drifting plankton, and is often numerous in nutrient-enriched stream segments.



CADDISFLIES


The most common beetles in streams are riffle beetles (adult and larva shown) and water pennies (not shown). Most of these require a swift current and an adequate supply of oxygen, and are generally considered clean-water indicators.



BEETLES


Appendix VII. B.
Aquatic Macroinvertebrates that Usually Indicate Poor Water Quality

Midges are the most common aquatic flies. The larvae occur in almost any aquatic situation. Many species are very tolerant to pollution. Large, red midge larvae called “bloodworms” indicate organic enrichment. Other midge larvae filter plankton, indicating nutrient enrichment when numerous.




MIDGES


Black fly larvae have specialized structures for filtering plankton and bacteria from the water, and require a strong current. Some species are tolerant of organic enrichment and toxic contaminants, while others are intolerant of pollutants.




BLACK FLIES



The segmented worms include the leeches and the small aquatic worms. The latter are more common, though usually unnoticed. They burrow in the substrate and feed on bacteria in the sediment. They can thrive under conditions of severe pollution and very low oxygen levels, and are thus valuable pollution indicators. Many leeches are also tolerant of poor water quality.




WORMS



Aquatic sowbugs are crustaceans that are often numerous in situations of high organic content and low oxygen levels. They are classic indicators of sewage pollution, and can also thrive in toxic situations.

Digital images by Larry Abele, New York State Department of Environmental Conservation, Stream Biomonitoring Unit.



SOWBUGS

Appendix VIII. The Rationale of Biological Monitoring

Biological monitoring refers to the use of resident benthic macroinvertebrate communities as indicators of water quality. Macroinvertebrates are larger-than-microscopic invertebrate animals that inhabit aquatic habitats; freshwater forms are primarily aquatic insects, worms, clams, snails, and crustaceans.

Concept

Nearly all streams are inhabited by a community of benthic macroinvertebrates. The species comprising the community each occupy a distinct niche defined and limited by a set of environmental requirements. The composition of the macroinvertebrate community is thus determined by many factors, including habitat, food source, flow regime, temperature, and water quality. The community is presumed to be controlled primarily by water quality if the other factors are determined to be constant or optimal. Community components which can change with water quality include species richness, diversity, balance, abundance, and presence/absence of tolerant or intolerant species. Various indices or metrics are used to measure these community changes. Assessments of water quality are based on metric values of the community, compared to expected metric values.

Advantages

The primary advantages of using macroinvertebrates as water quality indicators are that they:

- are sensitive to environmental impacts
- are less mobile than fish, and thus cannot avoid discharges
- can indicate effects of spills, intermittent discharges, and lapses in treatment
- are indicators of overall, integrated water quality, including synergistic effects
- are abundant in most streams and are relatively easy and inexpensive to sample
- are able to detect non-chemical impacts to the habitat, e.g. siltation or thermal changes
- are vital components of the aquatic ecosystem and important as a food source for fish
- are more readily perceived by the public as tangible indicators of water quality
- can often provide an on-site estimate of water quality
- can often be used to identify specific stresses or sources of impairment
- can be preserved and archived for decades, allowing for direct comparison of specimens
- bioaccumulate many contaminants, so that analysis of their tissues is a good monitor of toxic substances in the aquatic food chain

Limitations

Biological monitoring is not intended to replace chemical sampling, toxicity testing, or fish surveys. Each of these measurements provides information not contained in the others. Similarly, assessments based on biological sampling should not be taken as being representative of chemical sampling. Some substances may be present in levels exceeding ambient water quality criteria, yet have no apparent adverse community impact.

Appendix IX. Glossary

anthropogenic: caused by human actions

assessment: a diagnosis or evaluation of water quality

benthos: organisms occurring on or in the bottom substrate of a waterbody

bioaccumulate: accumulate contaminants in the tissues of an organism

biomonitoring: the use of biological indicators to measure water quality

community: a group of populations of organisms interacting in a habitat

drainage basin: an area in which all water drains to a particular waterbody; watershed

EPT richness: the number of species of mayflies (Ephemeroptera), stoneflies (Plecoptera), and caddisflies (Trichoptera) in a sample or subsample

facultative: occurring over a wide range of water quality; neither tolerant nor intolerant of poor water quality

fauna: the animal life of a particular habitat

impact: a change in the physical, chemical, or biological condition of a waterbody

impairment: a detrimental effect caused by an impact

index: a number, metric, or parameter derived from sample data used as a measure of water quality

intolerant: unable to survive poor water quality

ISC: Impervious Surface Cover; i.e., rooftops, roadways and parking areas - surfaces that cover soils which previously allowed rainwater infiltration.

longitudinal trends: upstream-downstream changes in water quality in a river or stream

macroinvertebrate: a larger-than-microscopic invertebrate animal that lives at least part of its life in aquatic habitats

multiplate: multiple-plate sampler, a type of artificial substrate sampler of aquatic macroinvertebrates

organism: a living individual

PAHs: Polycyclic Aromatic Hydrocarbons, a class of organic compounds that are often toxic or carcinogenic.

rapid bioassessment: a biological diagnosis of water quality using field and laboratory analysis designed to allow assessment of water quality in a short turn-around time; usually involves kick sampling and laboratory subsampling of the sample

riffle: wadeable stretch of stream usually with a rubble bottom and sufficient current to have the water surface broken by the flow; rapids

species richness: the number of macroinvertebrate species in a sample or subsample

station: a sampling site on a waterbody

survey: a set of samplings conducted in succession along a stretch of stream

synergistic effect: an effect produced by the combination of two factors that is greater than the sum of the two factors

tolerant: able to survive poor water quality

Appendix X. Impact Source Determination: Methods and Community Models

Definition Impact Source Determination (ISD) is the procedure for identifying types of impacts that exert deleterious effects on a waterbody. While the analysis of benthic macroinvertebrate communities has been shown to be an effective means of determining severity of water quality impacts, it has been less effective in determining what kind of pollution is causing the impact. ISD uses community types or models to ascertain the primary factor influencing the fauna.

Development of methods The method found to be most useful in differentiating impacts in New York State streams was the use of community types based on composition by family and genus. It may be seen as an elaboration of Percent Model Affinity (Novak and Bode, 1992), which is based on class and order. A large database of macroinvertebrate data was required to develop ISD methods. The database included several sites known or presumed to be impacted by specific impact types. The impact types were mostly known by chemical data or land use. These sites were grouped into the following general categories: agricultural nonpoint, toxic-stressed, sewage (domestic municipal), sewage/toxic, siltation, impoundment, and natural. Each group initially contained 20 sites. Cluster analysis was then performed within each group, using percent similarity at the family or genus level. Within each group four clusters were identified. Each cluster was usually composed of 4-5 sites with high biological similarity. From each cluster, a hypothetical model was then formed to represent a model cluster community type; sites within the cluster had at least 50 percent similarity to this model. These community type models formed the basis for ISD (see tables following). The method was tested by calculating percent similarity to all models and determining which model was the most similar to the test site. Some models were initially adjusted to achieve maximum representation of the impact type. New models are developed when similar communities are recognized from several streams.

Use of the ISD methods Impact Source Determination is based on similarity to existing models of community types (see tables following). The model that exhibits the highest similarity to the test data denotes the likely impact source type, or may indicate "natural," lacking an impact. In the graphic representation of ISD, only the highest similarity of each source type is identified. If no model exhibits a similarity to the test data of greater than 50 percent the determination is inconclusive. The determination of impact source type is used in conjunction with assessment of severity of water quality impact to provide an overall assessment of water quality.

Limitations These methods were developed for data derived from subsamples of 100-organisms each that are taken from traveling kick samples of New York State streams. Application of these methods for data derived from other sampling methods, habitats, or geographical areas would likely require modification of the models.

ISD MODELS TABLE
NATURAL MACROINVERTEBRATE COMMUNITY TYPE

	A	B	C	D	E	F	G	H	I	J	K	L	M
PLATYHELMINTHES	-	-	-	-	-	-	-	-	-	-	-	-	-
OLIGOCHAETA	-	-	5	-	5	-	5	5	-	-	-	5	5
HIRUDINEA	-	-	-	-	-	-	-	-	-	-	-	-	-
GASTROPODA	-	-	-	-	-	-	-	-	-	-	-	-	-
SPHAERIIDAE	-	-	-	-	-	-	-	-	-	-	-	-	-
ASELLIDAE	-	-	-	-	-	-	-	-	-	-	-	-	-
GAMMARIDAE	-	-	-	-	-	-	-	-	-	-	-	-	-
<u>Isonychia</u>	5	5	-	5	20	-	-	-	-	-	-	-	-
BAETIDAE	20	10	10	10	10	5	10	10	10	10	5	15	40
HEPTAGENIIDAE	5	10	5	20	10	5	5	5	5	10	10	5	5
LEPTOPHLEBIIDAE	5	5	-	-	-	-	-	-	5	-	-	25	5
EPHEMERELLIDAE	5	5	5	10	-	10	10	30	-	5	-	10	5
<u>Caenis/Tricorythodes</u>	-	-	-	-	-	-	-	-	-	-	-	-	-
PLECOPTERA	-	-	-	5	5	-	5	5	15	5	5	5	5
<u>Psephenus</u>	5	-	-	-	-	-	-	-	-	-	-	-	-
<u>Optioservus</u>	5	-	20	5	5	-	5	5	5	5	-	-	-
<u>Promoresia</u>	5	-	-	-	-	-	25	-	-	-	-	-	-
<u>Stenelmis</u>	10	5	10	10	5	-	-	-	10	-	-	-	5
PHILOPOTAMIDAE	5	20	5	5	5	5	5	-	5	5	5	5	5
HYDROPSYCHIDAE	10	5	15	15	10	10	5	5	10	15	5	5	10
HELICOPSYCHIDAE/ BRACHYCENTRIDAE/													
RHYACOPHILIDAE	5	5	-	-	-	20	-	5	5	5	5	5	-
SIMULIIDAE	-	-	-	5	5	-	-	-	-	5	-	-	-
<u>Simulium vittatum</u>	-	-	-	-	-	-	-	-	-	-	-	-	-
EMPIDIDAE	-	-	-	-	-	-	-	-	-	-	-	-	-
TIPULIDAE	-	-	-	-	-	-	-	-	5	-	-	-	-
CHIRONOMIDAE													
Tanypodinae	-	5	-	-	-	-	-	-	5	-	-	-	-
Diamesinae	-	-	-	-	-	-	5	-	-	-	-	-	-
Cardiocladius	-	5	-	-	-	-	-	-	-	-	-	-	-
<u>Cricotopus/</u>													
<u>Orthocladius</u>	5	5	-	-	10	-	-	5	-	-	5	5	5
<u>Eukiefferiella/</u>													
<u>Tvetenia</u>	5	5	10	-	-	5	5	5	-	5	-	5	5
<u>Parametriocnemus</u>	-	-	-	-	-	-	-	5	-	-	-	-	-
<u>Chironomus</u>	-	-	-	-	-	-	-	-	-	-	-	-	-
<u>Polypedilum aviceps</u> -	-	-	-	-	20	-	-	10	20	20	5	-	-
<u>Polypedilum</u> (all others)	5	5	5	5	5	-	5	5	-	-	-	-	-
Tanytarsini	-	5	10	5	5	20	10	10	10	10	40	5	5
TOTAL	100	100	100	100	100	100	100	100	100	100	100	100	100

ISD MODELS TABLE (cont.)
NONPOINT NUTRIENT ENRICHMENT IMPACTED MACROINVERTEBRATE COMMUNITY TYPE

	A	B	C	D	E	F	G	H	I	J
PLATYHELMINTHES	-	-	-	-	-	-	-	-	-	-
OLIGOCHAETA	-	-	-	5	-	-	-	-	-	15
HIRUDINEA	-	-	-	-	-	-	-	-	-	-
GASTROPODA	-	-	-	-	-	-	-	-	-	-
SPHAERIIDAE	-	-	-	5	-	-	-	-	-	-
ASELLIDAE	-	-	-	-	-	-	-	-	-	-
GAMMARIDAE	-	-	-	5	-	-	-	-	-	-
<u>Isonychia</u>	-	-	-	-	-	-	-	5	-	-
BAETIDAE	5	15	20	5	20	10	10	5	10	5
HEPTAGENIIDAE	-	-	-	-	5	5	5	5	-	5
LEPTOPHLEBIIDAE	-	-	-	-	-	-	-	-	-	-
EPHEMERELLIDAE	-	-	-	-	-	-	-	5	-	-
<u>Caenis/Tricorythodes</u>	-	-	-	-	5	-	-	5	-	5
PLECOPTERA	-	-	-	-	-	-	-	-	-	-
<u>Psephenus</u>	5	-	-	5	-	5	5	-	-	-
<u>Optioservus</u>	10	-	-	5	-	-	15	5	-	5
<u>Promoresia</u>	-	-	-	-	-	-	-	-	-	-
<u>Stenelmis</u>	15	15	-	10	15	5	25	5	10	5
PHILOPOTAMIDAE	15	5	10	5	-	25	5	-	-	-
HYDROPSYCHIDAE	15	15	15	25	10	35	20	45	20	10
HELICOPSYCHIDAE/ BRACHYCENTRIDAE/ RHYACOPHILIDAE	-	-	-	-	-	-	-	-	-	-
SIMULIIDAE	5	-	15	5	5	-	-	-	40	-
<u>Simulium vittatum</u>	-	-	-	-	-	-	-	-	5	-
EMPIDIDAE	-	-	-	-	-	-	-	-	-	-
TIPULIDAE	-	-	-	-	-	-	-	-	-	5
CHIRONOMIDAE										
Tanypodinae	-	-	-	-	-	-	5	-	-	5
<u>Cardiocladius</u>	-	-	-	-	-	-	-	-	-	-
<u>Cricotopus/</u> <u>Orthocladius</u>	10	15	10	5	-	-	-	-	5	5
<u>Eukiefferiella/</u> <u>Tvetenia</u>	-	15	10	5	-	-	-	-	5	-
<u>Parametriocnemus</u>	-	-	-	-	-	-	-	-	-	-
<u>Microtendipes</u>	-	-	-	-	-	-	-	-	-	20
<u>Polypedilum aviceps</u> -	-	-	-	-	-	-	-	-	-	-
<u>Polypedilum</u> (all others)	10	10	10	10	20	10	5	10	5	5
Tanytarsini	10	10	10	5	20	5	5	10	-	10
TOTAL	100	100	100	100	100	100	100	100	100	100

ISD MODELS TABLE (cont.)
 MACROINVERTEBRATE COMMUNITY TYPES
 MUNICIPAL/INDUSTRIAL WASTES IMPACTED TOXICS IMPACTED

	A	B	C	D	E	F	G	H	A	B	C	D	E	F
PLATYHELMINTHES	-	40	-	-	-	5	-	-	-	-	-	-	5	-
OLIGOCHAETA	20	20	70	10	-	20	-	-	-	10	20	5	5	15
HIRUDINEA	-	5	-	-	-	-	-	-	-	-	-	-	-	-
GASTROPODA	-	-	-	-	-	5	-	-	-	5	-	-	-	5
SPHAERIIDAE	-	5	-	-	-	-	-	-	-	-	-	-	-	-
ASELLIDAE	10	5	10	10	15	5	-	-	10	10	-	20	10	5
GAMMARIDAE	40	-	-	-	15	-	5	5	5	-	-	-	5	5
<u>Isonychia</u>	-	-	-	-	-	-	-	-	-	-	-	-	-	-
BAETIDAE	5	-	-	-	5	-	10	10	15	10	20	-	-	5
HEPTAGENIIDAE	5	-	-	-	-	-	-	-	-	-	-	-	-	-
LEPTOPHLEBIIDAE	-	-	-	-	-	-	-	-	-	-	-	-	-	-
EPHEMERELLIDAE	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<u>Caenis/Tricorythodes</u>	-	-	-	-	-	-	-	-	-	-	-	-	-	-
PLECOPTERA	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<u>Psephenus</u>	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<u>Criptoservus</u>	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<u>Promoresia</u>	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<u>Stenelmis</u>	5	-	-	10	5	-	5	5	10	15	-	40	35	5
PHILOPOTAMIDAE	-	-	-	-	-	-	-	40	10	-	-	-	-	-
HYDROPSYCHIDAE	10	-	-	50	20	-	40	20	20	10	15	10	35	10
HELICOPSYCHIDAE/ BRACHYCENTRIDAE/ RHYACOPHILIDAE	-	-	-	-	-	-	-	-	-	-	-	-	-	-
SIMULIIDAE	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<u>Simulium vittatum</u>	-	-	-	-	-	-	20	10	-	20	-	-	-	5
EMPIDIDAE	-	5	-	-	-	-	-	-	-	-	-	-	-	-
CHIRONOMIDAE	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Tanypodinae	-	10	-	-	5	15	-	-	5	10	-	-	-	25
<u>Cardiocladius</u>	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<u>Cricotopus/</u> <u>Orthocladius</u>	5	10	20	-	5	10	5	5	15	10	25	10	5	10
<u>Eukiefferiella/</u> <u>Tvetenia</u>	-	-	-	-	-	-	-	-	-	-	20	10	-	-
<u>Parametriocnemus</u>	-	-	-	-	-	-	-	-	-	-	-	5	-	-
<u>Chironomus</u>	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<u>Polypedilum aviceps</u>	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<u>Polypedilum</u> (all others)	-	-	-	10	20	40	10	5	10	-	-	-	-	5
Tanytarsini	-	-	-	10	10	-	5	-	-	-	-	-	-	5
TOTAL	100	100	100	100	100	100	100	100	100	100	100	100	100	100

ISD MODELS TABLE (cont.)
SEWAGE EFFLUENT, ANIMAL WASTES IMPACTED MACROINVERTEBRATE COMMUNITY TYPE

	A	B	C	D	E	F	G	H	I	J
PLATYHELMINTHES	-	-	-	-	-	-	-	-	-	-
OLIGOCHAETA	5	35	15	10	10	35	40	10	20	15
HIRUDINEA	-	-	-	-	-	-	-	-	-	-
GASTROPODA	-	-	-	-	-	-	-	-	-	-
SPHAERIIDAE	-	-	-	10	-	-	-	-	-	-
ASELLIDAE	5	10	-	10	10	10	10	50	-	5
GAMMARIDAE	-	-	-	-	-	10	-	10	-	-
<u>Isonychia</u>	-	-	-	-	-	-	-	-	-	-
BAETIDAE	-	10	10	5	-	-	-	-	5	-
HEPTAGENIIDAE	10	10	10	-	-	-	-	-	-	-
LEPTOPHLEBIIDAE	-	-	-	-	-	-	-	-	-	-
EPHEMERELLIDAE	-	-	-	-	-	-	-	-	5	-
<u>Caenis/Tricorythodes</u>	-	-	-	-	-	-	-	-	-	-
PLECOPTERA	-	-	-	-	-	-	-	-	-	-
<u>Psephenus</u>	-	-	-	-	-	-	-	-	-	-
<u>Optioservus</u>	-	-	-	-	-	-	-	-	5	-
<u>Promoresia</u>	-	-	-	-	-	-	-	-	-	-
<u>Stenelmis</u>	15	-	10	10	-	-	-	-	-	-
PHILOPOTAMIDAE	-	-	-	-	-	-	-	-	-	-
HYDROPSYCHIDAE	45	-	10	10	10	-	-	10	5	-
HELICOPSYCHIDAE/ BRACHYCENTRIDAE/ RHYACOPHILIDAE	-	-	-	-	-	-	-	-	-	-
SIMULIIDAE	-	-	-	-	-	-	-	-	-	-
<u>Simulium vittatum</u>	-	-	-	25	10	35	-	-	5	5
EMPIDIDAE	-	-	-	-	-	-	-	-	-	-
CHIRONOMIDAE	-	-	-	-	-	-	-	-	-	-
Tanypodinae	-	5	-	-	-	-	-	-	5	5
<u>Cardiocladius</u>	-	-	-	-	-	-	-	-	-	-
<u>Cricotopus/</u> <u>Orthocladius</u>	-	10	15	-	-	10	10	-	5	5
<u>Eukiefferiella/</u> <u>Tvetenia</u>	-	-	10	-	-	-	-	-	-	-
<u>Parametriocnemus</u>	-	-	-	-	-	-	-	-	-	-
<u>Chironomus</u>	-	-	-	-	-	-	10	-	-	60
<u>Polypedilum aviceps</u> -	-	-	-	-	-	-	-	-	-	-
<u>Polypedilum</u> (all others)	10	10	10	10	60	-	30	10	5	5
Tanytarsini	10	10	10	10	-	-	-	10	40	-
TOTAL	100	100	100	100	100	100	100	100	100	100

ISD MODELS TABLE (cont.)
MACROINVERTEBRATE COMMUNITY TYPES
SILTATION IMPACTED IMPOUNDMENT IMPACTED

	A	B	C	D	E	A	B	C	D	E	F	G	H	I	J
PLATYHELMINTHES	-	-	-	-	-	-	10	-	10	-	5	-	50	10	-
OLIGOCHAETA	5	-	20	10	5	5	-	40	5	10	5	10	5	5	-
HIRUDINEA	-	-	-	-	-	-	-	-	-	5	-	-	-	-	-
GASTROPODA	-	-	-	-	-	-	-	10	-	5	5	-	-	-	-
SPHAERIIDAE	-	-	-	5	-	-	-	-	-	-	-	-	5	25	-
ASELLIDAE	-	-	-	-	-	-	5	5	-	10	5	5	5	-	-
GAMMARIDAE	-	-	-	10	-	-	-	10	-	10	50	-	5	10	-
<u>Isonychia</u>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
BAETIDAE	-	10	20	5	-	-	5	-	5	-	-	5	-	-	5
HEPTAGENIIDAE	5	10	-	20	5	5	5	-	5	5	5	5	-	5	5
LEPTOPHLEBIIDAE	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
EPHEMERELLIDAE	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<u>Caenis/Tricorythodes</u>	5	20	10	5	15	-	-	-	-	-	-	-	-	-	-
PLECOPTERA	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<u>Psephenus</u>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	5
<u>Optioservus</u>	5	10	-	-	-	-	-	-	-	-	-	-	-	5	-
<u>Promoresia</u>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<u>Stenelmis</u>	5	10	10	5	20	5	5	10	10	-	5	35	-	5	10
PHILOPOTAMIDAE	-	-	-	-	-	5	-	-	5	-	-	-	-	-	30
HYDROPSYCHIDAE	25	10	-	20	30	50	15	10	10	10	10	20	5	15	20
HELICOPSYCHIDAE/ BRACHYCENTRIDAE/ RHYACOPHILIDAE	-	-	-	-	-	-	-	-	-	-	-	-	-	5	-
SIMULIIDAE	5	10	-	-	5	5	-	5	-	35	10	5	-	-	15
EMPIDIDAE	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
CHIRONOMIDAE															
Tanypodinae	-	-	-	-	-	-	5	-	-	-	-	-	-	-	-
<u>Cardiocladius</u>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<u>Cricotopus/</u> <u>Orthocladius</u>	25	-	10	5	5	5	25	5	-	10	-	5	10	-	-
<u>Eukiefferiella/</u> <u>Tvetenia</u>	-	-	10	-	5	5	15	-	-	-	-	-	-	-	-
<u>Parametriocnemus</u>	-	-	-	-	-	5	-	-	-	-	-	-	-	-	-
<u>Chironomus</u>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<u>Polypedilum aviceps</u> - <u>Polypedilum</u> (all others)	10	10	10	5	5	5	-	-	20	-	-	5	5	5	5
Tanytarsini	10	10	10	10	5	5	10	5	30	-	-	5	10	10	5
TOTAL	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100

APPENDIX XI. METHODS FOR CALCULATING OF THE NUTRIENT BIOTIC INDEX

Definition: The Nutrient Biotic Index (Smith, 2005) is a diagnostic measure of stream nutrient enrichment identified by macroinvertebrate taxa. The frequency of occurrences of taxa at varying nutrient concentrations allowed the identification of taxon-specific nutrient optima using a method of weighted averaging. The establishment of nutrient optima is possible based on the observation that most species exhibit unimodal response curves in relation to environmental variables (Jongman et al. 1987). The assignment of tolerance values to taxa based on their nutrient optimum provided the ability to reduce macroinvertebrate community data to a linear scale of eutrophication from oligotrophic to eutrophic. Two tolerance values were assigned to each taxon, one for total phosphorus, and one for nitrate (listed in Smith, 2005). This provides the ability to calculate two different nutrient biotic indices, one for total phosphorus (NBI-P), and one for nitrate (NBI-N). Study of the indices indicate better performance by the NBI-P, with strong correlations to stream nutrient status assessment based on diatom information.

Calculation of the NBI-P and NBI-N: Calculation of the indices [2] follows the approach of Hilsenhoff (1987).

$$\text{NBI Score}_{(\text{TP or NO}_3^-)} = \sum (a \times b) / c$$

Where *a* is equal to the number of individuals for each taxon, *b* is the taxon's tolerance value, and *c* is the total number of individuals in the sample for which tolerance values have been assigned.

Classification of NBI Scores: NBI scores have been placed on a scale of eutrophication with provisional boundaries between stream trophic status.

Index	Oligotrophic	Mesotrophic	Eutrophic
NBI-P	< 5.0	> 5.0 - 6.5	> 6.5
NBI-N	< 4.5	> 4.5 - 6.0	> 6.0

References:

- Hilsenhoff, W. L., 1987, An improved biotic index of organic stream pollution. *The Great Lakes Entomologist* 20(1): 31-39.
- Jongman, R. H. G., C. J. F. ter Braak and O. F. R. van Tongeren, 1987, *Data analysis in community and landscape ecology*. Pudoc Wageningen, Netherlands, 299 pages.
- Smith, A.J., 2005, *Development of a Nutrient Biotic Index for use with benthic macroinvertebrates*. Masters Thesis, SUNY Albany. 70 pages.

Appendix XII. Characteristics of Headwater Stream Sites

Headwater stream sites are defined as first or second-order locations close to the source, usually less than three miles. Natural characteristics of headwaters sometimes result in erroneous assessment of water quality.

The following are typical characteristics of headwater sites:

- **Reduced upstream community recruitment populations** reduce drift colonization, and may reduce species richness.
- **Usually nutrient-poor, lower in food resources, and less productive.**
- **A few intolerant species may be very abundant**, due to reduced, simplified fauna. For a 100-organism subsample, this can affect species richness, EPT richness, and percent model affinity. The dominant species averages 37% of the total fauna, and is an intolerant species of either mayfly (e.g., Epeorus, Paraleptophlebia, Stenonema), stonefly (e.g., Leuctridae or Capniidae), caddisfly (e.g., Brachycentrus, Dolophilodes, or Chimarra), or riffle beetle (e.g., Optioservus or Promoresia).
- **Many community indices are low**, even though invertebrate communities are dominated by intolerant species. Average index values are: species richness - 19, EPT richness - 8, Hilsenhoff biotic index - 3.05, and percent model affinity - 57 (based on headwaters of a number of New York State streams).

Due to the above characteristics, it is recommended that corrective action be taken to adjust for non-representative indices from headwater sites. *A correction factor of 1.5 may be applied to species richness, EPT richness, and percent model affinity.* Criteria for the use of the correction factor are: a headwater location as described above; a community dominated by an intolerant species, and species richness, EPT richness, and percent model affinity are judged to be non-representative of actual water quality. *Alternatively, index values may be maintained, and the overall assessment may be adjusted up to non-impacted if the above criteria are met.*

APPENDIX XIII. Calculation of Impervious Surface Cover Using Orthoimagery.

All data development and analysis are conducted using ArcGIS ArcView 9.1.

1. Delineation of watershed and site-location subbasin boundaries. Either of two methods may be used; the second of these was used for calculation in the Quackenderry Creek watershed.
 - a. Digital boundary delineation using the 10m Digital Elevation Models (DEM) and NYS-DEC Hydrography Network, and Waterbodies layer in ArcGIS. This method uses an automated procedure developed by the Martyn J. Smith of the USGS, 425 Jordan Road, Troy, NY 12180 (marSmith@usgs.gov).
 - b. Delineation by hand using a hardcopy version of *USGS 7.5' Topographic Quadrangle* and with the resulting boundaries approved by a USGS staff hydrologist. Watershed boundaries are then "heads-up digitized" (traced by hand using mouse) as a GIS layer using a digital version of the same quadrangle obtained from the Cornell University Geospatial Information Repository (URL: <http://cugir.mannlib.cornell.edu/>).
2. Calculation of the area of each subbasin: use the Geo-processing Tools within ArcGIS.
3. Delineation of Impervious Surfaces: Impervious surfaces - including paved surfaces (roads, parking lots, driveways etc.), buildings, pools, paths, and walkways - are heads-up digitized into a GIS layer, using the latest 12-inch resolution, natural color orthoimagery from the NYS Office of Cyber Security & Critical Infrastructure Coordination. Twenty-four-inch resolution, color-infrared orthoimagery is used when available, to confirm the existence or absence of impervious surfaces in areas of dense vegetation or shadow present in the natural-color imagery. Both data sets can be obtained from the New York State GIS Clearinghouse (URL: <http://www.nysgis.state.ny.us/>).
4. Calculation of Percent Impervious Surface for Each Subbasin: The impervious surface layer is overlaid with the subbasin layer in ArcGIS. Geo-processing is then used to create a new layer, which subsequently allows for the derivation of the impervious surface areas within each subbasin. The sum of the impervious surface areas is then divided by the area of the entire subbasin and multiplied by 100 to calculate percent impervious surface area of the subbasin.

