

102 10/10



New York State  
**DEPARTMENT OF ENVIRONMENTAL CONSERVATION**

STR  
500-2  
P0003  
204-4413

Division of Water

# Mourning Kill

## Biological Assessment

2005 Survey



New York State  
**Department of Environmental Conservation**

George E. Pataki, Governor

Denise M. Sheehan, Commissioner





# New York State Documents

OCLC:



CHECKLIST: May 2006

Original Document Scanned at:

- 400 DPI       Simplex  
 Duplex

Original Document contained:

- Black & White Photos  
 Colored Photos  
 Colored Print (list color) \_\_\_\_\_  
 Colored Paper (list color) \_\_\_\_\_  
 Line Art, Graphs  
 Oversized Pages -- reduced from \_\_\_\_\_ (original size)  
 Text Only

Date Scanned: 8/13/2008

**This electronic document has been scanned by the  
New York State Library from a paper original and has been stored  
on optical media.**

**The New York State Library  
Cultural Education Center  
Albany, NY 12230**

(MASTER.DOC. 9/99)



# **Mourning Kill**


## BIOLOGICAL ASSESSMENT

Upper Hudson River Basin  
Saratoga County, New York

Survey date: September 21, 2005  
Report date: March 9, 2006

Robert W. Bode  
Margaret A. Novak  
Lawrence E. Abele  
Diana L. Heitzman  
Alexander J. Smith

Stream Biomonitoring Unit  
Bureau of Water Assessment and Management  
Division of Water  
NYS Department of Environmental Conservation  
Albany, New York




## CONTENTS

Background.....	1
Results and Conclusions.....	1
Discussion.....	2
Literature Cited.....	3
Overview of Field Data.....	3
Figure 1. Biological Assessment Profile, 2005.....	4
Table 1. Impact Source Determination.....	5
Table 2. Station Locations.....	6
Figure 2. Site Overview Map .....	7
Figures 3a-c. Site Location Maps.....	8
Table 3. Macroinvertebrate Species Collected.....	11
Macroinvertebrate Data Reports: Raw Data .....	12
Laboratory Data Summary.....	16
Field Data Summary.....	17
Appendix I. Biological Methods for Kick Sampling.....	18
Appendix II. Macroinvertebrate Community Parameters.....	19
Appendix III. Levels of Water Quality Impact in Streams.....	20
Appendix IV. Biological Assessment Profile Derivation.....	21
Appendix V. Water Quality Assessment Criteria.....	23
Appendix VI. Traveling Kick Sample Illustration.....	24
Appendix VII. Macroinvertebrate Illustrations.....	25
Appendix VIII. Rationale for Biological Monitoring.....	27
Appendix IX. Glossary.....	28
Appendix X. Impact Source Determination: Methods and Community Models .....	29
Appendix XI. Nutrient Biotic Index .....	35

NEW YORK STATE LIBRARY



B00464489B



**Stream:** Mourning Kill, Saratoga County, New York

**Reach:** Harmony Corners to Ballston Spa, New York

**NYS Drainage Basin:** Upper Hudson River

### **Background:**

The Stream Biomonitoring Unit sampled the Mourning Kill in Saratoga County, New York, on September 21, 2005. The purpose of the sampling was to assess overall water quality and establish a baseline of data from which to compare future results.

In riffle areas at four sites, a traveling kick sample for macroinvertebrates was taken using methods described in the Quality Assurance document (Bode et al., 2002) and summarized in Appendix I. The contents of each sample were field-inspected to determine major groups of organisms present, and then preserved in alcohol for laboratory inspection of a 100-specimen subsample from each site. Macroinvertebrate community parameters used in the determination of water quality included species richness, biotic index, EPT richness, and percent model affinity (see Appendices II and III). Expected variability of results is stated in Smith and Bode (2004). Table 2 provides a listing of sampling sites and Table 3 provides a listing of all macroinvertebrate species collected in the present survey. This is followed by macroinvertebrate data reports, including raw data from each site.

### **Results and Conclusions:**

1. Water quality in the Mourning Kill was assessed as slightly impacted at all sites, based on resident macroinvertebrate communities.
2. As indicators of development in the watershed, future studies of the Mourning Kill should particularly monitor the parameters of conductivity and the Nutrient Biotic Index. The stonefly species *Agneta capitata* could also be monitored as a clean-water indicator whose continued presence reflects acceptable water quality in the stream.

## Discussion:

The Mourning Kill begins near O'Brien Corners in Saratoga County, New York, and flows approximately 17 miles in an easterly direction before flowing into Kayaderosseras Creek at Ballston Spa. The stream is classified as C, meaning the best water use is for fishing and fish propagation. The stream has not been previously sampled by the Stream Biomonitoring Unit. The purpose of the present sampling was to assess overall water quality and establish a baseline of data from which to compare future results.

In the present study water quality in the Mourning Kill was assessed as slightly impacted at all four sites (Figure 1). No longitudinal trends in water quality were discerned from source to mouth. Resident macroinvertebrate communities at all sites were dominated by caddisflies, mayflies and riffle beetles. All sites contained clean-water stoneflies, but overall, species richness was poor. The Impact Source Determination (ISD) of *complex* at Stations-2 and -4 (Table 1) is not explained by any known discharges, and is likely spurious. Community Model H in the *complex* category is dominated by caddisflies of the genus *Philopotamidae*, and is similar to Community Model J in the *impoundment* category (see Appendix X). ISD values are high in both the *complex* and *impoundment* categories at Stations-2 and -4. Sluggish wetland-like conditions in some reaches of the Mourning Kill may simulate impoundment effects in benthic macroinvertebrates, resulting in the *Philopotamidae*-dominated communities.

The Nutrient Biotic Index (NBI), a metric recently developed by Smith (2005) to evaluate levels of nutrient enrichment, is included in the Biological Assessment Profile in this report (Appendix XI). Overall water quality assessments are thus based on the average of five metrics (see Appendix II). Values for NBI-P (total phosphorus) in the Mourning Kill ranged from 4.55 to 5.60. Values for NBI-N (nitrates) ranged from 5.01 to 5.56. Both were below the level of eutrophic conditions (6.0), indicating that nutrient enrichment is not a controlling factor for macroinvertebrate communities.

Future studies of the Mourning Kill should target the parameters of conductivity and nutrients as indicators of development in the watershed. The stonefly *Agnatina capitata* could be monitored as a clean-water indicator whose continued presence reflects acceptable water quality in the stream. The less sensitive stonefly *Paragnetina media*, which was found at all sites, would be expected to survive slight impacts that may eliminate *Agnatina capitata*.

## **Literature Cited:**

Bode, R. W., M. A. Novak, L. E. Abele, D. L. Heitzman and A. J. Smith, 2002, Quality assurance work plan for biological stream monitoring in New York State. New York State Department of Environmental Conservation, Technical Report, 115 pages.

Smith, A.J., 2005, Development of a Nutrient Biotic Index for use with benthic macroinvertebrates. SUNY Albany, Masters Thesis, 70 pages.

Smith, A. J. and R. W. Bode, 2004, Analysis of variability in New York State benthic macroinvertebrate samples. New York State Department of Environmental Conservation, Technical Report, 43 pages.

## **Overview of field data:**

On September 21, 2005, the Mourning Kill at the sites sampled was 2-6 meters wide, 0.1-0.2 meters deep, and had current speeds of 40-90 cm/sec in riffles. Dissolved oxygen was 8.3- 9.9 mg/l, specific conductance was 422-522  $\mu$ mhos, pH was 7.6-8.2 and temperature was 15.0-18.0 °C (59-64 °F). Measurements for each site are found on the field data summary sheets.

Figure 1. Biological Assessment Profile of index values, Mourning Kill, 2005. Values are plotted on a normalized scale of water quality. The line connects the mean of five values for each site, representing species richness, EPT richness, Hilsenhoff Biotic Index, Nutrient Biotic Index (P) and Percent Model Affinity. See Appendix IV for a more complete explanation.




Table 1. Impact Source Determination, Mourning Kill, 2005. Numbers represent percent similarity to community type models for each impact category. Highest similarities at each station are highlighted. Similarities less than 50% are less conclusive. Highest numbers represent probable type of impact. See Appendix X for further explanation.

Community Type	STATION			
	MORN-1	MORN-2	MORN-3	MORN-4
Natural: minimal human impacts	57	46	54	38
Nutrient additions: mostly nonpoint, agricultural	47	50	50	37
Toxic: industrial, municipal, or urban run-off	30	33	28	24
Organic: sewage effluent, animal wastes	27	30	27	16
Complex: municipal/industrial	37	59	25	45
Siltation	38	38	37	24
Impoundment	47	59	36	47

STATION      COMMUNITY TYPE  
MORN-1      Natural  
MORN-2      Complex, impoundment (see Discussion)  
MORN-3      Natural  
MORN-4      Complex, impoundment (see Discussion)

TABLE 2. Station Locations for the Mourning Kill, Saratoga County, NY

STATION      LOCATION

01      Harmony Corners, New York  
Below Route 67 bridge  
Latitude/Longitude 42° 59' 07"; 73° 55' 53"  
12.0 stream miles above mouth



02      Ballston Center, New York  
Above Goode Street bridge  
Latitude/Longitude 42° 57' 15"; 73° 54' 10"  
7.9 stream miles above mouth






03      Ballston Center, New York  
Above Middle Line Road bridge  
Latitude/Longitude 42° 58' 12"; 73° 52' 47"  
6.0 stream miles above mouth




04      Ballston Spa, New York  
Above Route 67 bridge  
Latitude/Longitude 42° 58' 33"; 73° 50' 54"  
3.9 stream miles above mouth












TABLE 3. Macroinvertebrate Species Collected in the Mourning Kill, Saratoga County, New York, 2005.

PLATYHELMINTHES	Helicopsychidae
TURBELLARIA	<i>Helicopsyche borealis</i>
Undetermined Turbellaria	DIPTERA
ARTHROPODA	Tipulidae
INSECTA	<i>Dicranota sp.</i>
EPHEMEROPTERA	Athericidae
Isonychiidae	<i>Atherix sp.</i>
<i>Isonychia bicolor</i>	Simuliidae
Baetidae	<i>Simulium tuberosum</i>
<i>Acentrella sp.</i>	Chironomidae
<i>Baetis flavistriga</i>	<i>Thienemannimyia gr. spp.</i>
<i>Baetis intercalaris</i>	<i>Nanocladius sp.</i>
Heptageniidae	<i>Parachaetocladius sp.</i>
<i>Stenacron interpunctatum</i>	<i>Tvetenia bavarica gr.</i>
<i>Stenonema sp.</i>	<i>Polypedilum aviceps</i>
Ephemerellidae	<i>Tanytarsus guerlus gr.</i>
Undetermined Ephemerellidae	
Leptophlebiidae	
Undetermined Leptophlebiidae	
PLECOPTERA	
Perlidae	
<i>Agnatina capitata</i>	
<i>Paragnetina media</i>	
COLEOPTERA	
Psephenidae	
<i>Psephenus herricki</i>	
Elmidae	
<i>Optioservus fastiditus</i>	
<i>Optioservus sp.</i>	
<i>Stenelmis crenata</i>	
<i>Stenelmis sp.</i>	
MEGALOPTERA	
Corydalidae	
<i>Nigronia serricornis</i>	
Sialidae	
<i>Sialis sp.</i>	
TRICHOPTERA	
Philopotamidae	
<i>Chimarra aterrima?</i>	
Hydropsychidae	
<i>Cheumatopsyche sp.</i>	
<i>Hydropsyche bronta</i>	
<i>Hydropsyche slossonae</i>	
Rhyacophilidae	
<i>Rhyacophila sp.</i>	
Hydroptilidae	
<i>Leucotrichia sp.</i>	

## Macroinvertebrate Data Reports: Raw Data

STREAM SITE: Mourning Kill, Station MORN-01  
 LOCATION: Harmony Corners, NY, below Route 67 bridge  
 DATE: 21 September 2005  
 SAMPLE TYPE: Kick sample  
 SUBSAMPLE: 100 organisms

### PLATYHELMINTHES

TURBELLARIA	Planariidae	Undetermined Turbellaria	2
-------------	-------------	--------------------------	---

### ARTHROPODA

#### INSECTA

EPHEMEROPTERA	Isonychiidae	<i>Isonychia bicolor</i>	6
	Baetidae	<i>Baetis intercalaris</i>	1
	Heptageniidae	<i>Stenonema sp.</i>	22
	Leptophlebiidae	Undetermined Leptophlebiidae	3
PLECOPTERA	Perlidae	<i>Agnetina capitata</i>	4
COLEOPTERA	Psephenidae	<i>Psephenus herricki</i>	9
	Elmidae	<i>Optioservus sp.</i>	5
		<i>Stenelmis sp.</i>	3
MEGALOPTERA	Sialidae	<i>Sialis sp.</i>	1
TRICHOPTERA	Philopotamidae	<i>Chimarra aterrima?</i>	19
	Hydropsychidae	<i>Cheumatopsyche sp.</i>	13
		<i>Hydropsyche bronta</i>	1
DIPTERA	Tipulidae	<i>Dicranota sp.</i>	4
	Chironomidae	<i>Thienemannimyia gr. spp.</i>	2
		<i>Parachaetocladius sp.</i>	3
		<i>Tvetenia bavarica gr.</i>	1
		<i>Tanytarsus guerlus gr.</i>	1

SPECIES RICHNESS: 18 (poor)  
 BIOTIC INDEX: 3.77 (very good)  
 EPT RICHNESS: 8 (good)  
 MODEL AFFINITY: 70 (very good)  
 ASSESSMENT: slightly impacted (6.88)  
 NUTRIENT BI, NBI-P 5.30 (mesotrophic)

DESCRIPTION: The kick sample was taken 20 meters downstream of the Route 67 bridge. The stream was only 2 meters wide, and current speed was slow. The macroinvertebrate community appeared well-balanced, with clean-water mayflies, stoneflies, caddisflies, beetles and hellgrammites, but species richness was low, likely due to headwater status. Based on the metrics, water quality was assessed as slightly impacted.

## Macroinvertebrate Data Reports: Raw Data, cont'd.

STREAM SITE: Mourning Kill, Station MORN- 02  
 LOCATION: Ballston Center, NY, above Goode Street bridge  
 DATE: 21 September 2005  
 SAMPLE TYPE: Kick sample  
 SUBSAMPLE: 100 organisms

### ARTHROPODA

#### INSECTA

EPHEMEROPTERA	Baetidae	<i>Acentrella sp.</i>	1
	Heptageniidae	<i>Stenonema sp.</i>	8
	Ephemerellidae	Undetermined Ephemerellidae	1
PLECOPTERA	Perlidae	<i>Agetina capitata</i>	1
		<i>Paragnetina media</i>	2
COLEOPTERA	Elmidae	<i>Optioservus fastiditus</i>	9
		<i>Stenelmis crenata</i>	9
MEGALOPTERA	Corydalidae	<i>Nigronia serricornis</i>	8
TRICHOPTERA	Philopotamidae	<i>Chimarra aterrima?</i>	40
	Hydropsychidae	<i>Cheumatopsyche sp.</i>	3
<i>Hydropsyche bronta</i>		7	
<i>Hydropsyche slossonae</i>		3	
DIPTERA	Tipulidae	<i>Dicranota sp.</i>	1
		<i>Nanocladius sp.</i>	1
		<i>Parachaetocladius sp.</i>	5
		<i>Polypedilum aviceps</i>	1

SPECIES RICHNESS: 16 (poor)  
 BIOTIC INDEX: 4.06 (very good)  
 EPT RICHNESS: 9 (good)  
 MODEL AFFINITY: 49 (poor)  
 ASSESSMENT: slightly impacted (5.99)  
 NUTRIENT BI. NBI-P 4.63 (oligotrophic)

DESCRIPTION: This site had a higher current speed than Station-1 and appeared to be an ideal macroinvertebrate habitat. Mayflies, stoneflies, and hellgrammites were present, but the community was heavily dominated by caddisflies. Based on the metrics, water quality was assessed as slightly impacted.

Macroinvertebrate Data Reports: Raw Data, cont'd.

STREAM SITE: Mourning Kill, Station MORN-03  
 LOCATION: Ballston Center, NY, above Middle Line Road bridge  
 DATE: 21 September 2005  
 SAMPLE TYPE: Kick sample  
 SUBSAMPLE: 100 organisms

ARTHROPODA

INSECTA

EPHEMEROPTERA	Isonychiidae	<i>Isonychia bicolor</i>	8
	Baetidae	<i>Baetis flavistriga</i>	1
		<i>Baetis intercalaris</i>	5
	Heptageniidae	<i>Stenacron interpunctatum</i>	1
		<i>Stenonema sp.</i>	20
PLECOPTERA	Perlidae	<i>Paragnetina media</i>	2
COLEOPTERA	Psephenidae	<i>Psephenus herricki</i>	2
	Elmidae	<i>Optioservus fastiditus</i>	22
		<i>Stenelmis crenata</i>	15
MEGALOPTERA	Corydalidae	<i>Nigronia serricornis</i>	4
TRICHOPTERA	Philopotamidae	<i>Chimarra aterrima?</i>	13
	Hydropsychidae	<i>Hydropsyche slossonae</i>	1
	Rhyacophilidae	<i>Rhyacophila sp.</i>	1
DIPTERA	Athericidae	<i>Atherix sp.</i>	2
	Chironomidae	<i>Thienemannimyia gr. spp.</i>	1
		<i>Nanocladius sp.</i>	2

SPECIES RICHNESS: 16 (poor)  
 BIOTIC INDEX: 3.92 (very good)  
 EPT RICHNESS: 9 (good)  
 MODEL AFFINITY: 66 (very good)  
 ASSESSMENT: slightly impacted (6.71)  
 NUTRIENT BI, NBI-P 5.60 (mesotrophic)

DESCRIPTION: The sampling site was 30 meters upstream of the Middle Line Road bridge. Habitat was adequate for macroinvertebrates, but the stream bottom had more silt than at Station-2. The community was dominated by algal-scraping riffle beetles. Based on the metrics, water quality was assessed as slightly impacted.

Macroinvertebrate Data Reports: Raw Data, cont'd.

STREAM SITE: Mourning Kill, Station MORN- 04  
 LOCATION: Ballston Spa, NY, above Route 67 bridge  
 DATE: 21 September 2005  
 SAMPLE TYPE: Kick sample  
 SUBSAMPLE: 100 organisms

ARTHROPODA

INSECTA

EPHEMEROPTERA	Isonychiidae	<i>Isonychia bicolor</i>	14
	Heptageniidae	<i>Stenonema sp.</i>	2
PLECOPTERA	Perlidae	<i>Paragnetina media</i>	2
COLEOPTERA	Elmidae	<i>Optioservus sp.</i>	7
		<i>Stenelmis crenata</i>	10
MEGALOPTERA	Corydalidae	<i>Nigronia serricornis</i>	5
TRICHOPTERA	Philopotamidae	<i>Chimarra aterrima?</i>	36
	Hydropsychidae	<i>Cheumatopsyche sp.</i>	1
		<i>Hydropsyche bronta</i>	3
	Hydroptilidae	<i>Leucotrichia sp.</i>	1
	Helicopsychidae	<i>Helicopsyche borealis</i>	1
DIPTERA	Tipulidae	<i>Dicranota sp.</i>	14
	Simuliidae	<i>Simulium tuberosum</i>	1
	Athericidae	<i>Atherix sp.</i>	1
	Chironomidae	<i>Parachaetocladius sp.</i>	2

SPECIES RICHNESS: 15 (poor)  
 BIOTIC INDEX: 3.70 (very good)  
 EPT RICHNESS: 8 (good)  
 MODEL AFFINITY: 50 (good)  
 ASSESSMENT: slightly impacted (5.95)  
 NUTRIENT BI, NBI-P 4.55 (oligotrophic)

DESCRIPTION: The kick sample was taken approximately 60 meters upstream of Route 67, upstream of the railroad trestle. Brown algae was noticed in the stream that had not been observed at upstream sites. The macroinvertebrate community was dominated by caddisflies. Based on the metrics, water quality was assessed as slightly impacted.

## LABORATORY DATA SUMMARY

<b>STREAM NAME: Mourning Kill</b>		<b>DRAINAGE: 11</b>		
<b>DATE SAMPLED: 9/21/2005</b>		<b>COUNTY: Saratoga</b>		
<b>SAMPLING METHOD: Travelling Kick</b>				
<b>STATION</b>	01	02	03	04
<b>LOCATION</b>	Harmony Corners	Ballston Center	Ballston Center	Ballston Spa
<b>DOMINANT SPECIES/% CONTRIBUTION/TOLERANCE/COMMON NAME</b>				
1.	<i>Stenonema sp.</i>	<i>Chimarra aterrima?</i>	<i>Optioservus fastiditus</i>	<i>Chimarra aterrima?</i>
	22 %	40 %	22 %	36 %
	intolerant	intolerant	intolerant	intolerant
	mayfly	caddisfly	beetle	caddisfly
2.	<i>Chimarra aterrima?</i>	<i>Optioservus fastiditus</i>	<i>Stenonema sp.</i>	<i>Isonychia bicolor</i>
<b>Intolerant = not tolerant of poor water quality</b>	19 %	9 %	20 %	14 %
	intolerant	intolerant	intolerant	intolerant
	caddisfly	beetle	mayfly	mayfly
3.	<i>Cheumatopsyche sp.</i>	<i>Stenelmis crenata</i>	<i>Stenelmis crenata</i>	<i>Dicranota sp.</i>
<b>Facultative = occurring over a wide range of water quality</b>	13 %	9 %	15 %	14 %
	facultative	facultative	facultative	intolerant
	caddisfly	beetle	beetle	crane fly
4.	<i>Psephenus herricki</i>	<i>Stenonema sp.</i>	<i>Chimarra aterrima?</i>	<i>Stenelmis crenata</i>
<b>Tolerant = tolerant of poor water quality</b>	9 %	8 %	13 %	10 %
	facultative	intolerant	intolerant	facultative
	beetle	mayfly	caddisfly	beetle
5.	<i>Isonychia bicolor</i>	<i>Nigronia serricornis</i>	<i>Isonychia bicolor</i>	<i>Optioservus sp.</i>
	6 %	8 %	8 %	7 %
	intolerant	facultative	intolerant	intolerant
	mayfly	megaloptera	mayfly	beetle
<b>% CONTRIBUTION OF MAJOR GROUPS (NUMBER OF TAXA IN PARENTHESES)</b>				
<b>Chironomidae (midges)</b>	7.0 (4.0)	7.0 (3.0)	3.0 (2.0)	2.0 (1.0)
<b>Trichoptera (caddisflies)</b>	33.0 (3.0)	53.0 (4.0)	15.0 (3.0)	42.0 (5.0)
<b>Ephemeroptera (mayflies)</b>	32.0 (4.0)	10.0 (3.0)	35.0 (5.0)	16.0 (2.0)
<b>Plecoptera (stoneflies)</b>	4.0 (1.0)	3.0 (2.0)	2.0 (1.0)	2.0 (1.0)
<b>Coleoptera (beetles)</b>	17.0 (3.0)	18.0 (2.0)	39.0 (3.0)	17.0 (2.0)
<b>Oligochaeta (worms)</b>	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)
<b>Mollusca (clams and snails)</b>	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)
<b>Crustacea (crayfish, scuds, sowbugs)</b>	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)
<b>Other insects (odonates, diptera)</b>	5.0 (2.0)	9.0 (2.0)	6.0 (2.0)	21.0 (4.0)
<b>Other (Nemertea, Platyhelminthes)</b>	2.0 (1.0)	0.0 (0.0)	0.0 (0.0)	1.0 (1.0)
<b>SPECIES RICHNESS</b>	18	16	16	15
<b>BIOTIC INDEX</b>	3.77	4.06	3.92	3.70
<b>EPT RICHNESS</b>	8	9	9	8
<b>PERCENT MODEL AFFINITY</b>	70	49	66	50
<b>FIELD ASSESSMENT</b>	Very good	Very good	Very good	Very good
<b>OVERALL ASSESSMENT</b>	Slight	Slight	Slight	Slight

## FIELD DATA SUMMARY

<b>STREAM NAME: Mourning Kill</b>		<b>DATE SAMPLED: 9/21/2005</b>		
<b>REACH: Harmony Corners - Ballston Spa</b>				
<b>FIELD PERSONNEL INVOLVED: Bode, Smith</b>				
<b>STATION</b>	01	02	03	04
<b>ARRIVAL TIME AT STATION</b>	09:30 AM	10:10 AM	10:45 AM	11:10 AM
<b>LOCATION</b>	Harmony Corners	Ballston Center	Ballston Center	Ballston Spa
<b>PHYSICAL CHARACTERISTICS</b>				
Width (meters)	2.0	4.0	6.0	6.0
Depth (meters)	0.1	0.1	0.2	0.1
Current speed (cm per sec.)	40	90	80	80
Substrate (%)				
Rock (>25.4 cm, or bedrock)				
Rubble (6.35 – 25.4 cm)	40	40	40	40
Gravel (0.2 – 6.35 cm)	30	30	30	30
Sand (0.06 – 2.0 mm)	10	20	10	10
Silt (0.004 – 0.06 mm)	20	10	20	20
Embeddedness (%)	20	30	30	30
<b>CHEMICAL MEASUREMENTS</b>				
Temperature (° C)	16.0	15.0	17.0	18.0
Specific Conductance (umhos)	422	494	511	522
Dissolved Oxygen (mg/l)	9.2	8.3	9.9	8.9
pH	7.6	7.9	7.9	8.2
<b>BIOLOGICAL ATTRIBUTES</b>				
Canopy (%)	60	95	50	80
Aquatic Vegetation				
algae – suspended				
algae – attached, filamentous				
algae – diatoms	x	x	x	x
macrophytes or moss				
Occurrence of Macroinvertebrates				
Ephemeroptera (mayflies)	x	x	x	x
Plecoptera (stoneflies)	x	x	x	x
Trichoptera (caddisflies)	x	x	x	x
Coleoptera (beetles)	x	x	x	x
Megaloptera (dobsonflies, alderflies)	x	x	x	x
Odonata (dragonflies, damselflies)				
Chironomidae (midges)				x
Simuliidae (black flies)				x
Decapoda (crayfish)	x	x	x	x
Gammaridae (scuds)				
Mollusca (snails, clams)				
Oligochaeta (worms)				
Other		x		x
<b>FAUNAL CONDITION</b>	Very good	Very good	Very good	Very good

## Appendix I. Biological Methods for Kick Sampling

A. Rationale: The use of the standardized kick sampling method provides a biological assessment technique that lends itself to rapid assessments of stream water quality.

B. Site Selection: Sampling sites are selected based on these criteria: (1) The sampling location should be a riffle with a substrate of rubble, gravel and sand; depth should be one meter or less, and current speed should be at least 0.4 meters per second. (2) The site should have comparable current speed, substrate type, embeddedness, and canopy cover to both upstream and downstream sites to the degree possible. (3) Sites are chosen to have a safe and convenient access.

C. Sampling: Macroinvertebrates are sampled using the standardized traveling kick method. An aquatic net is positioned in the water at arms' length downstream and the stream bottom is disturbed by foot, so that organisms are dislodged and carried into the net. Sampling is continued for a specified time and distance in the stream. Rapid assessment sampling specifies sampling for five minutes over a distance of five meters. The contents of the net are emptied into a pan of stream water. The contents are then examined, and the major groups of organisms are recorded, usually on the ordinal level (e.g., stoneflies, mayflies, caddisflies). Larger rocks, sticks, and plants may be removed from the sample if organisms are first removed from them. The contents of the pan are poured into a U.S. No. 30 sieve and transferred to a quart jar. The sample is then preserved by adding 95% ethyl alcohol.

D. Sample Sorting and Subsampling: In the laboratory, the sample is rinsed with tap water in a U.S. No. 40 standard sieve to remove any fine particles left in the residues from field sieving. The sample is transferred to an enamel pan and distributed homogeneously over the bottom of the pan. A small amount of the sample is randomly removed with a spatula, rinsed with water, and placed in a petri dish. This portion is examined under a dissecting stereomicroscope and 100 organisms are randomly removed from the debris. As they are removed, they are sorted into major groups, placed in vials containing 70 percent alcohol, and counted. The total number of organisms in the sample is estimated by weighing the residue from the picked subsample and determining its proportion of the total sample weight.

E. Organism Identification: All organisms are identified to the species level whenever possible. Chironomids and oligochaetes are slide-mounted and viewed through a compound microscope; most other organisms are identified as whole specimens using a dissecting stereomicroscope. The number of individuals in each species and the total number of individuals in the subsample are recorded on a data sheet. All organisms from the subsample are archived (either slide-mounted or preserved in alcohol). If the results of the identification process are ambiguous, suspected of being spurious, or do not yield a clear water quality assessment, additional subsampling may be required.

## Appendix II. Macroinvertebrate Community Parameters

1. Species Richness: the total number of species or taxa found in the sample. For subsamples of 100-organisms each that are taken from kick samples, expected ranges in most New York State streams are: greater than 26, non-impacted; 19-26, slightly impacted; 11-18, moderately impacted; less than 11, severely impacted.
2. EPT Richness: the total number of species of mayflies (Ephemeroptera), stoneflies (Plecoptera), and caddisflies (Trichoptera) found in an average 100-organisms subsample. These are considered to be clean-water organisms, and their presence is generally correlated with good water quality (Lenat, 1987). Expected assessment ranges from most New York State streams are: greater than 10, non-impacted; 6-10, slightly impacted; 2-5, moderately impacted; and 0-1, severely impacted.
3. Hilsenhoff Biotic Index: a measure of the tolerance of organisms in a sample to organic pollution (sewage effluent, animal wastes) and low dissolved oxygen levels. It is calculated by multiplying the number of individuals of each species by its assigned tolerance value, summing these products, and dividing by the total number of individuals. On a 0-10 scale, tolerance values range from intolerant (0) to tolerant (10). For the purpose of characterizing species' tolerance, intolerant = 0-4, facultative = 5-7, and tolerant = 8-10. Tolerance values are listed in Hilsenhoff (1987). Additional values are assigned by the NYS Stream Biomonitoring Unit. The most recent values for each species are listed in Quality Assurance document, Bode et al. (1996). Impact ranges are: 0-4.50, non-impacted; 4.51-6.50, slightly impacted; 6.51-8.50, moderately impacted; and 8.51-10.00, severely impacted.
4. Percent Model Affinity: a measure of similarity to a model, non-impacted community based on percent abundance in seven major macroinvertebrate groups (Novak and Bode, 1992). Percentage abundances in the model community are: 40% Ephemeroptera; 5% Plecoptera; 10% Trichoptera; 10% Coleoptera; 20% Chironomidae; 5% Oligochaeta; and 10% Other. Impact ranges are: greater than 64, non-impacted; 50-64, slightly impacted; 35-49, moderately impacted; and less than 35, severely impacted.
5. Nutrient Biotic Index: a measure of stream nutrient enrichment identified by macroinvertebrate taxa. It is calculated by multiplying the number of individuals of each species by its assigned tolerance value, summing these products, and dividing by the total number of individuals with assigned tolerance values. Tolerance values ranging from intolerant (0) to tolerant (10) are based on nutrient optima for Total Phosphorus (listed in Smith, 2005). Impact ranges are: 0-5.00, non-impacted; 5.01-6.00, slightly impacted; 6.01-7.00, moderately impacted and 7.01-10.00, severely impacted.

### Appendix III. Levels of Water Quality Impact in Streams

The description of overall stream water quality based on biological parameters uses a four-tiered system of classification. Level of impact is assessed for each individual parameter and then combined for all parameters to form a consensus determination. Four parameters are used: species richness, EPT richness, biotic index, and percent model affinity (see Appendix II). The consensus is based on the determination of the majority of the parameters. Since parameters measure different aspects of the macroinvertebrate community, they cannot be expected to always form unanimous assessments. The assessment ranges given for each parameter are based on subsamples of 100-organisms each that are taken from macroinvertebrate riffle kick samples. These assessments also apply to most multiplate samples, with the exception of percent model affinity.

1. *Non-impacted*: Indices reflect very good water quality. The macroinvertebrate community is diverse, usually with at least 27 species in riffle habitats. Mayflies, stoneflies, and caddisflies are well represented; EPT richness is greater than 10. The biotic index value is 4.50 or less. Percent model affinity is greater than 64. Nutrient Biotic Index is 5.00 or less. Water quality should not be limiting to fish survival or propagation. This level of water quality includes both pristine habitats and those receiving discharges which minimally alter the biota.


2. *Slightly impacted*: Indices reflect good water quality. The macroinvertebrate community is slightly but significantly altered from the pristine state. Species richness is usually 19-26. Mayflies and stoneflies may be restricted, with EPT richness values of 6-10. The biotic index value is 4.51-6.50. Percent model affinity is 50-64. Nutrient Biotic Index is 5.01-6.00. Water quality is usually not limiting to fish survival, but may be limiting to fish propagation.

3. *Moderately impacted*: Indices reflect poor water quality. The macroinvertebrate community is altered to a large degree from the pristine state. Species richness is usually 11-18 species. Mayflies and stoneflies are rare or absent, and caddisflies are often restricted; the EPT richness is 2-5. The biotic index value is 6.51-8.50. Percent model affinity is 35-49. Nutrient Biotic Index is 6.01-7.00. Water quality often is limiting to fish propagation, but usually not to fish survival.

4. *Severely impacted*: Indices reflect very poor water quality. The macroinvertebrate community is limited to a few tolerant species. Species richness is 10 or fewer. Mayflies, stoneflies and caddisflies are rare or absent; EPT richness is 0-1. The biotic index value is greater than 8.50. Percent model affinity is less than 35. Nutrient Biotic Index is greater than 7.00. The dominant species are almost all tolerant, and are usually midges and worms. Often, 1-2 species are very abundant. Water quality is often limiting to both fish propagation and fish survival.

Appendix IV-A: Biological Assessment Profile (BAP); Conversion of Index Values to a Common 10-Scale

The Biological Assessment Profile of index values, developed by Phil O'Brien, Division of Water, NYSDEC, is a method of plotting biological index values on a common scale of water quality impact. Values from the five indices -- species richness (SPP), EPT richness (EPT), Hilsenhoff Biotic Index (HBI), Percent Model Affinity (PMA), and Nutrient Biotic Index (NBI)-- defined in Appendix II are converted to a common 0-10 scale using the formulae in the Quality Assurance document (Bode, et al., 2002), and as shown in the figure below.



## Appendix IV-B. Biological Assessment Profile: Plotting Values


To plot survey data:

1. Position each site on the x-axis according to miles or tenths of a mile upstream of the mouth.
2. Plot the values of the four indices for each site as indicated by the common scale.
3. Calculate the mean of the four values and plot the result. This represents the assessed impact for each site.

Example data:

	Station 1		Station 2	
	metric value	10-scale value	metric value	10-scale value
Species richness	20	5.59	33	9.44
Hilsenhoff Biotic Index	5.00	7.40	4.00	8.00
EPT richness	9	6.80	13	.00
Percent Model Affinity	55	5.97	65	7.60
Nutrient Biotic Index	6.50	3.75	3.50	9.00
Average		5.90 (slight)		8.61 (non-)

Sample Plot of Biological Assessment Profile values



Appendix V. Water Quality Assessment Criteria

Non-Navigable Flowing Waters

	Species Richness	Hilsenhoff Biotic Index	Nutrient Biotic Index*	EPT Value	Percent Model Affinity#	Diversity**
Non-Impacted	>26	0.00-4.50	0.00-5.00	>10	>64	>4
Slightly Impacted	19-26	4.51-6.50	5.01-6.00	6-10	50-64	3.01-4.00
Moderately Impacted	11-18	6.51-8.50	6.51-7.00	2-5	35-49	2.01-3.00
Severely Impacted	0-10	8.51-10.00	7.01-10.00	0-1	<35	0.00-2.00

\* Nutrient Biotic Index (for total phosphorus, NBI-P) used for traveling kick samples but not for multiplate samples.


# Percent model affinity criteria used for traveling kick samples but not for multiplate samples.

\*\* Diversity criteria are used for multiplate samples but not for traveling kick samples.

Navigable Flowing Waters

	Species	Hilsenhoff	EPT	Species
Non-	>21	0.00-7.00	>5	>3.00
Slightly	17-21	7.01-8.00	4-5	2.51-3.00
Moderately	12-16	8.01-9.00	2-3	2.01-2.50
Severely	0-11	9.01-10.00	0-1	0.00-2.00

## Appendix VI: The Traveling Kick Sample




← current

Rocks and sediment in a riffle are dislodged by foot upstream of a net. Dislodged organisms are carried by the current into the net. Sampling continues for five minutes, as the sampler gradually moves downstream to cover a distance of five meters.


## Appendix VII. A. Aquatic Macroinvertebrates that Usually Indicate Good Water Quality

Mayfly nymphs are often the most numerous organisms found in clean streams. They are sensitive to most types of pollution, including low dissolved oxygen (less than 5 ppm), chlorine, ammonia, metals, pesticides, and acidity. Most mayflies are found clinging to the undersides of rocks.




*MAYFLIES*

Stonefly nymphs are mostly limited to cool, well-oxygenated streams. They are sensitive to most of the same pollutants as mayflies, except acidity. They are usually much less numerous than mayflies. The presence of even a few stoneflies in a stream suggests that good water quality has been maintained for several months.




*STONEFLIES*

Caddisfly larvae often build a portable case of sand, stones, sticks, or other debris. Many caddisfly larvae are sensitive to pollution, although a few are tolerant. One family spins nets to catch drifting plankton, and is often numerous in nutrient-enriched stream segments.




*CADDISFLIES*

The most common beetles in streams are riffle beetles (adult and larva pictured) and water pennies (not shown). Most of these require a swift current and an adequate supply of oxygen, and are generally considered clean-water indicators.




*BEETLES*




Appendix VII. B. Aquatic Macroinvertebrates that Usually Indicate Poor Water Quality

Midges are the most common aquatic flies. The larvae occur in almost any aquatic situation. Many species are very tolerant to pollution. Large, red midge larvae called “bloodworms” indicate organic enrichment. Other midge larvae filter plankton, indicating nutrient enrichment when numerous.




*MIDGES*


Black fly larvae have specialized structures for filtering plankton and bacteria from the water, and require a strong current. Some species are tolerant of organic enrichment and toxic contaminants, while others are intolerant of pollutants.




*BLACK FLIES*




The segmented worms include the leeches and the small aquatic worms. The latter are more common, though usually unnoticed. They burrow in the substrate and feed on bacteria in the sediment. They can thrive under conditions of severe pollution and very low oxygen levels, and are thus valuable pollution indicators. Many leeches are also tolerant of poor water quality.



*WORMS*



Aquatic sowbugs are crustaceans that are often numerous in situations of high organic content and low oxygen levels. They are classic indicators of sewage pollution, and can also thrive in toxic situations.



*SOWBUGS*

Digital images by Larry Abele, New York State Department of Environmental Conservation, Stream Biomonitoring Unit.

## Appendix VIII. The Rationale of Biological Monitoring

Biological monitoring refers to the use of resident benthic macroinvertebrate communities as indicators of water quality. Macroinvertebrates are larger-than-microscopic invertebrate animals that inhabit aquatic habitats; freshwater forms are primarily aquatic insects, worms, clams, snails, and crustaceans.

### Concept:

Nearly all streams are inhabited by a community of benthic macroinvertebrates. The species comprising the community each occupy a distinct niche defined and limited by a set of environmental requirements. The composition of the macroinvertebrate community is thus determined by many factors, including habitat, food source, flow regime, temperature, and water quality. The community is presumed to be controlled primarily by water quality if the other factors are determined to be constant or optimal. Community components which can change with water quality include species richness, diversity, balance, abundance, and presence/absence of tolerant or intolerant species. Various indices or metrics are used to measure these community changes. Assessments of water quality are based on metric values of the community, compared to expected metric values.

### Advantages:

The primary advantages to using macroinvertebrates as water quality indicators are that they:

- are sensitive to environmental impacts
- are less mobile than fish, and thus cannot avoid discharges
- can indicate effects of spills, intermittent discharges, and lapses in treatment
- are indicators of overall, integrated water quality, including synergistic effects
- are abundant in most streams and are relatively easy and inexpensive to sample
- are able to detect non-chemical impacts to the habitat, e.g. siltation or thermal changes
- are vital components of the aquatic ecosystem and important as a food source for fish
- are more readily perceived by the public as tangible indicators of water quality
- can often provide an on-site estimate of water quality
- can often be used to identify specific stresses or sources of impairment
- can be preserved and archived for decades, allowing for direct comparison of specimens
- bioaccumulate many contaminants, so that analysis of their tissues is a good monitor of toxic substances in the aquatic food chain

### Limitations:

Biological monitoring is not intended to replace chemical sampling, toxicity testing, or fish surveys. Each of these measurements provides information not contained in the others. Similarly, assessments based on biological sampling should not be taken as being representative of chemical sampling. Some substances may be present in levels exceeding ambient water quality criteria, yet have no apparent adverse community impact.

## Appendix IX: Glossary

- anthropogenic:** caused by human actions
- assessment:** a diagnosis or evaluation of water quality
- benthos:** organisms occurring on or in the bottom substrate of a waterbody
- bioaccumulate:** accumulate contaminants in the tissues of an organism
- biomonitoring:** the use of biological indicators to measure water quality
- community:** a group of populations of organisms interacting in a habitat
- drainage basin:** an area in which all water drains to a particular waterbody; watershed
- electrofishing:** sampling fish by using electric currents to temporarily immobilize them, allowing capture
- EPT richness:** the number of species of mayflies (Ephemeroptera), stoneflies (Plecoptera), and caddisflies (Trichoptera) in a sample or subsample
- facultative:** occurring over a wide range of water quality; neither tolerant nor intolerant of poor water quality
- fauna:** the animal life of a particular habitat
- impact:** a change in the physical, chemical, or biological condition of a waterbody
- impairment:** a detrimental effect caused by an impact
- index:** a number, metric, or parameter derived from sample data used as a measure of water quality
- intolerant:** unable to survive poor water quality
- longitudinal trends:** upstream-downstream changes in water quality in a river or stream
- macroinvertebrate:** a larger-than-microscopic invertebrate animal that lives at least part of its life in aquatic habitats
- multiplate:** multiple-plate sampler, a type of artificial substrate sampler of aquatic macroinvertebrates
- organism:** a living individual
- PAHs:** Polycyclic Aromatic Hydrocarbons, a class of organic compounds that are often toxic or carcinogenic.
- rapid bioassessment:** a biological diagnosis of water quality using field and laboratory analysis designed to allow assessment of water quality in a short turn-around time; usually involves kick sampling and laboratory subsampling of the sample
- riffle:** wadeable stretch of stream usually with a rubble bottom and sufficient current to have the water surface broken by the flow; rapids
- species richness:** the number of macroinvertebrate species in a sample or subsample
- station:** a sampling site on a waterbody
- survey:** a set of samplings conducted in succession along a stretch of stream
- synergistic effect:** an effect produced by the combination of two factors that is greater than the sum of the two factors
- tolerant:** able to survive poor water quality

## Appendix X. Impact Source Determination Methods and Community Models

**Definition:** Impact Source Determination (ISD) is the procedure for identifying types of impacts that exert deleterious effects on a waterbody. While the analysis of benthic macroinvertebrate communities has been shown to be an effective means of determining severity of water quality impacts, it has been less effective in determining what kind of pollution is causing the impact. ISD uses community types or models to ascertain the primary factor influencing the fauna.

**Development of methods:** The method found to be most useful in differentiating impacts in New York State streams was the use of community types based on composition by family and genus. It may be seen as an elaboration of Percent Model Affinity (Novak and Bode, 1992), which is based on class and order. A large database of macroinvertebrate data was required to develop ISD methods. The database included several sites known or presumed to be impacted by specific impact types. The impact types were mostly known by chemical data or land use. These sites were grouped into the following general categories: agricultural nonpoint, toxic-stressed, sewage (domestic municipal), sewage/toxic, siltation, impoundment, and natural. Each group initially contained 20 sites. Cluster analysis was then performed within each group, using percent similarity at the family or genus level. Within each group, four clusters were identified. Each cluster was usually composed of 4-5 sites with high biological similarity. From each cluster, a hypothetical model was then formed to represent a model cluster community type; sites within the cluster had at least 50 percent similarity to this model. These community type models formed the basis for ISD (see tables following). The method was tested by calculating percent similarity to all the models and determining which model was the most similar to the test site. Some models were initially adjusted to achieve maximum representation of the impact type. New models are developed when similar communities are recognized from several streams.

**Use of the ISD methods:** Impact Source Determination is based on similarity to existing models of community types (see tables following). The model that exhibits the highest similarity to the test data denotes the likely impact source type, or may indicate "natural," lacking an impact. In the graphic representation of ISD, only the highest similarity of each source type is identified. If no model exhibits a similarity to the test data of greater than 50 percent, the determination is inconclusive. The determination of impact source type is used in conjunction with assessment of severity of water quality impact to provide an overall assessment of water quality.

**Limitations:** These methods were developed for data derived from subsamples of 100-organisms each that are taken from traveling kick samples of New York State streams. Application of these methods for data derived from other sampling methods, habitats, or geographical areas would likely require modification of the models.

ISD MODELS TABLE  
MACROINVERTEBRATE COMMUNITY TYPE  
NATURAL

	A	B	C	D	E	F	G	H	I	J	K	L	M
PLATYHELMINTHES	-	-	-	-	-	-	-	-	-	-	-	-	-
OLIGOCHAETA	-	-	5	-	5	-	5	5	-	-	-	5	5
HIRUDINEA	-	-	-	-	-	-	-	-	-	-	-	-	-
GASTROPODA	-	-	-	-	-	-	-	-	-	-	-	-	-
SPHAERIIDAE	-	-	-	-	-	-	-	-	-	-	-	-	-
ASELLIDAE	-	-	-	-	-	-	-	-	-	-	-	-	-
GAMMARIDAE	-	-	-	-	-	-	-	-	-	-	-	-	-
<u>Isonychia</u>	5	5	-	5	20	-	-	-	-	-	-	-	-
BAETIDAE	20	10	10	10	10	5	10	10	10	10	5	15	40
HEPTAGENIIDAE	5	10	5	20	10	5	5	5	5	10	10	5	5
LEPTOPHLEBIIDAE	5	5	-	-	-	-	-	-	5	-	-	25	5
EPHEMERELLIDAE	5	5	5	10	-	10	10	30	-	5	-	10	5
<u>Caenis/Tricorythodes</u>	-	-	-	-	-	-	-	-	-	-	-	-	-
PLECOPTERA	-	-	-	5	5	-	5	5	15	5	5	5	5
<u>Psephenus</u>	5	-	-	-	-	-	-	-	-	-	-	-	-
<u>Optioservus</u>	5	-	20	5	5	-	5	5	5	5	-	-	-
<u>Promoresia</u>	5	-	-	-	-	-	25	-	-	-	-	-	-
<u>Stenelmis</u>	10	5	10	10	5	-	-	-	10	-	-	-	5
PHILOPOTAMIDAE	5	20	5	5	5	5	5	-	5	5	5	5	5
HYDROPSYCHIDAE	10	5	15	15	10	10	5	5	10	15	5	5	10
HELICOPSYCHIDAE/ BRACHYCENTRIDAE/ RHYACOPHILIDAE	5	5	-	-	-	20	-	5	5	5	5	5	-
SIMULIIDAE	-	-	-	5	5	-	-	-	-	5	-	-	-
<u>Simulium vittatum</u>	-	-	-	-	-	-	-	-	-	-	-	-	-
EMPIDIDAE	-	-	-	-	-	-	-	-	-	-	-	-	-
TIPULIDAE	-	-	-	-	-	-	-	-	5	-	-	-	-
CHIRONOMIDAE													
Tanyptodinae	-	5	-	-	-	-	-	-	5	-	-	-	-
Diamesinae	-	-	-	-	-	-	5	-	-	-	-	-	-
Cardiocladius	-	5	-	-	-	-	-	-	-	-	-	-	-
<u>Cricotopus/</u> <u>Orthocladius</u>	5	5	-	-	10	-	-	5	-	-	5	5	5
<u>Eukiefferiella/</u> <u>Tvetenia</u>	5	5	10	-	-	5	5	5	-	5	-	5	5
<u>Parametricnemus</u>	-	-	-	-	-	-	-	5	-	-	-	-	-
<u>Chironomus</u>	-	-	-	-	-	-	-	-	-	-	-	-	-
<u>Polypedilum aviceps</u> -	-	-	-	-	20	-	-	10	20	20	5	-	-
<u>Polypedilum</u> (all others)	5	5	5	5	5	-	5	5	-	-	-	-	-
Tanytarsini	-	5	10	5	5	20	10	10	10	10	40	5	5
TOTAL	100	100	100	100	100	100	100	100	100	100	100	100	100

ISD MODELS TABLE (cont.)  
 MACROINVERTEBRATE COMMUNITY TYPE  
 NONPOINT NUTRIENT ENRICHMENT IMPACTED

	A	B	C	D	E	F	G	H	I	J
PLATYHELMINTHES	-	-	-	-	-	-	-	-	-	-
OLIGOCHAETA	-	-	-	5	-	-	-	-	-	15
HIRUDINEA	-	-	-	-	-	-	-	-	-	-
GASTROPODA	-	-	-	-	-	-	-	-	-	-
SPHAERIIDAE	-	-	-	5	-	-	-	-	-	-
ASELLIDAE	-	-	-	-	-	-	-	-	-	-
GAMMARIDAE	-	-	-	5	-	-	-	-	-	-
<u>Isonychia</u>	-	-	-	-	-	-	-	5	-	-
BAETIDAE	5	15	20	5	20	10	10	5	10	5
HEPTAGENIIDAE	-	-	-	-	5	5	5	5	-	5
LEPTOPHLEBIIDAE	-	-	-	-	-	-	-	-	-	-
EPHEMERELLIDAE	-	-	-	-	-	-	-	5	-	-
<u>Caenis/Tricorythodes</u>	-	-	-	-	5	-	-	5	-	5
PLECOPTERA	-	-	-	-	-	-	-	-	-	-
<u>Psephenus</u>	5	-	-	5	-	5	5	-	-	-
<u>Optioservus</u>	10	-	-	5	-	-	15	5	-	5
<u>Promoesia</u>	-	-	-	-	-	-	-	-	-	-
<u>Stenelmis</u>	15	15	-	10	15	5	25	5	10	5
PHILOPOTAMIDAE	15	5	10	5	-	25	5	-	-	-
HYDROPSYCHIDAE	15	15	15	25	10	35	20	45	20	10
HELICOPSYCHIDAE/ BRACHYCENTRIDAE/ RHYACOPHILIDAE	-	-	-	-	-	-	-	-	-	-
SIMULIIDAE	5	-	15	5	5	-	-	-	40	-
<u>Simulium vittatum</u>	-	-	-	-	-	-	-	-	5	-
EMPIDIDAE	-	-	-	-	-	-	-	-	-	-
TIPULIDAE	-	-	-	-	-	-	-	-	-	5
CHIRONOMIDAE										
Tanypodinae	-	-	-	-	-	-	5	-	-	5
<u>Cardiocladius</u>	-	-	-	-	-	-	-	-	-	-
<u>Cricotopus/</u> <u>Orthocladius</u>	10	15	10	5	-	-	-	-	5	5
<u>Eukiefferiella/</u> <u>Tvetenia</u>	-	15	10	5	-	-	-	-	5	-
<u>Parametriocnemus</u>	-	-	-	-	-	-	-	-	-	-
<u>Microtendipes</u>	-	-	-	-	-	-	-	-	-	20
<u>Polypedilum aviceps</u>	-	-	-	-	-	-	-	-	-	-
<u>Polypedilum</u> (all others)	10	10	10	10	20	10	5	10	5	5
Tanytarsini	10	10	10	5	20	5	5	10	-	10
TOTAL	100	100	100	100	100	100	100	100	100	100

ISD MODELS TABLE (cont.)  
 MACROINVERTEBRATE COMMUNITY TYPES  
 MUNICIPAL/INDUSTRIAL WASTES IMPACTED                      TOXICS IMPACTED

	A	B	C	D	E	F	G	H	A	B	C	D	E	F
PLATYHELMINTHES	-	40	-	-	-	5	-	-	-	-	-	-	5	-
OLIGOCHAETA	20	20	70	10	-	20	-	-	-	10	20	5	5	15
HIRUDINEA	-	5	-	-	-	-	-	-	-	-	-	-	-	-
GASTROPODA	-	-	-	-	-	5	-	-	-	5	-	-	-	5
SPHAERIIDAE	-	5	-	-	-	-	-	-	-	-	-	-	-	-
ASELLIDAE	10	5	10	10	15	5	-	-	10	10	-	20	10	5
GAMMARIDAE	40	-	-	-	15	-	5	5	5	-	-	-	5	5
<u>Isonychia</u>	-	-	-	-	-	-	-	-	-	-	-	-	-	-
BAETIDAE	5	-	-	-	5	-	10	10	15	10	20	-	-	5
HEPTAGENIIDAE	5	-	-	-	-	-	-	-	-	-	-	-	-	-
LEPTOPHLEBIIDAE	-	-	-	-	-	-	-	-	-	-	-	-	-	-
EPHEMERELLIDAE	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<u>Caenis/Tricorythodes</u>	-	-	-	-	-	-	-	-	-	-	-	-	-	-
PLECOPTERA	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<u>Psephenus</u>	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<u>Optioservus</u>	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<u>Promoresia</u>	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<u>Stenelmis</u>	5	-	-	10	5	-	5	5	10	15	-	40	35	5
PHILOPOTAMIDAE	-	-	-	-	-	-	-	40	10	-	-	-	-	-
HYDROPSYCHIDAE	10	-	-	50	20	-	40	20	20	10	15	10	35	10
HELICOPSYCHIDAE/ BRACHYCENTRIDAE/ RHYACOPHILIDAE	-	-	-	-	-	-	-	-	-	-	-	-	-	-
SIMULIIDAE	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<u>Simulium vittatum</u>	-	-	-	-	-	-	20	10	-	20	-	-	-	5
EMPIDIDAE	-	5	-	-	-	-	-	-	-	-	-	-	-	-
CHIRONOMIDAE														
Tanypodinae	-	10	-	-	5	15	-	-	5	10	-	-	-	25
<u>Cardiocladius</u>	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<u>Cricotopus/</u> <u>Orthocladius</u>	5	10	20	-	5	10	5	5	15	10	25	10	5	10
<u>Eukiefferiella/</u> <u>Tvetenia</u>	-	-	-	-	-	-	-	-	-	-	20	10	-	-
<u>Parametriocnemus</u>	-	-	-	-	-	-	-	-	-	-	-	5	-	-
<u>Chironomus</u>	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<u>Polypedilum aviceps</u>	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<u>Polypedilum</u> (all others)	-	-	-	10	20	40	10	5	10	-	-	-	-	5
Tanytarsini	-	-	-	10	10	-	5	-	-	-	-	-	-	5
TOTAL	100	100	100	100	100	100	100	100	100	100	100	100	100	100

ISD MODELS TABLE (cont.)  
 MACROINVERTEBRATE COMMUNITY TYPE  
 SEWAGE EFFLUENT, ANIMAL WASTES IMPACTED

	A	B	C	D	E	F	G	H	I	J
PLATYHELMINTHES	-	-	-	-	-	-	-	-	-	-
OLIGOCHAETA	5	35	15	10	10	35	40	10	20	15
HIRUDINEA	-	-	-	-	-	-	-	-	-	-
GASTROPODA	-	-	-	-	-	-	-	-	-	-
SPHAERIIDAE	-	-	-	10	-	-	-	-	-	-
ASELLIDAE	5	10	-	10	10	10	10	50	-	5
GAMMARIDAE	-	-	-	-	-	10	-	10	-	-
<u>Isonychia</u>	-	-	-	-	-	-	-	-	-	-
BAETIDAE	-	10	10	5	-	-	-	-	5	-
HEPTAGENIIDAE	10	10	10	-	-	-	-	-	-	-
LEPTOPHLEBIIDAE	-	-	-	-	-	-	-	-	-	-
EPHEMERELLIDAE	-	-	-	-	-	-	-	-	5	-
<u>Caenis/Tricorythodes</u>	-	-	-	-	-	-	-	-	-	-
PLECOPTERA	-	-	-	-	-	-	-	-	-	-
<u>Psephenus</u>	-	-	-	-	-	-	-	-	-	-
<u>Optioservus</u>	-	-	-	-	-	-	-	-	5	-
<u>Promoresia</u>	-	-	-	-	-	-	-	-	-	-
<u>Stenelmis</u>	15	-	10	10	-	-	-	-	-	-
PHILOPOTAMIDAE	-	-	-	-	-	-	-	-	-	-
HYDROPSYCHIDAE	45	-	10	10	10	-	-	10	5	-
HELICOPSYCHIDAE/ BRACHYCENTRIDAE/ RHYACOPHILIDAE	-	-	-	-	-	-	-	-	-	-
SIMULIIDAE	-	-	-	-	-	-	-	-	-	-
<u>Simulium vittatum</u>	-	-	-	25	10	35	-	-	5	5
EMPIDIDAE	-	-	-	-	-	-	-	-	-	-
CHIRONOMIDAE	-	-	-	-	-	-	-	-	-	-
Tanypodinae	-	5	-	-	-	-	-	-	5	5
<u>Cardiocladius</u>	-	-	-	-	-	-	-	-	-	-
<u>Cricotopus/</u> <u>Orthocladius</u>	-	10	15	-	-	10	10	-	5	5
<u>Eukiefferiella/</u> <u>Tvetenia</u>	-	-	10	-	-	-	-	-	-	-
<u>Parametrioconemus</u>	-	-	-	-	-	-	-	-	-	-
<u>Chironomus</u>	-	-	-	-	-	-	10	-	-	60
<u>Polypedilum aviceps</u>	-	-	-	-	-	-	-	-	-	-
<u>Polypedilum</u> (all others)	10	10	10	10	60	-	30	10	5	5
Tanytarsini	10	10	10	10	-	-	-	10	40	-
TOTAL	100	100	100	100	100	100	100	100	100	100

ISD MODELS TABLE (cont.)  
 MACROINVERTEBRATE COMMUNITY TYPES  
 SILTATION IMPACTED                      IMPOUNDMENT IMPACTED

	A	B	C	D	E	A	B	C	D	E	F	G	H	I	J
PLATYHELMINTHES	-	-	-	-	-	-	10	-	10	-	5	-	50	10	-
OLIGOCHAETA	5	-	20	10	5	5	-	40	5	10	5	10	5	5	-
HIRUDINEA	-	-	-	-	-	-	-	-	-	5	-	-	-	-	-
GASTROPODA	-	-	-	-	-	-	-	10	-	5	5	-	-	-	-
SPHAERIIDAE	-	-	-	5	-	-	-	-	-	-	-	-	5	25	-
ASELLIDAE	-	-	-	-	-	-	5	5	-	10	5	5	5	-	-
GAMMARIDAE	-	-	-	10	-	-	-	10	-	10	50	-	5	10	-
<u>Isonychia</u>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
BAETIDAE	-	10	20	5	-	-	5	-	5	-	-	5	-	-	5
HEPTAGENIIDAE	5	10	-	20	5	5	5	-	5	5	5	5	-	5	5
LEPTOPHLEBIIDAE	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
EPHEMERELLIDAE	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<u>Caenis/Tricorythodes</u>	5	20	10	5	15	-	-	-	-	-	-	-	-	-	-
PLECOPTERA	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<u>Psephenus</u>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	5
<u>Optioservus</u>	5	10	-	-	-	-	-	-	-	-	-	-	-	5	-
<u>Promoresia</u>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<u>Stenelmis</u>	5	10	10	5	20	5	5	10	10	-	5	35	-	5	10
PHILOPOTAMIDAE	-	-	-	-	-	5	-	-	5	-	-	-	-	-	30
HYDROPSYCHIDAE	25	10	-	20	30	50	15	10	10	10	10	20	5	15	20
HELICOPSYCHIDAE/ BRACHYCENTRIDAE/ RHYACOPHILIDAE	-	-	-	-	-	-	-	-	-	-	-	-	-	5	-
SIMULIIDAE	5	10	-	-	5	5	-	5	-	35	10	5	-	-	15
EMPIDIDAE	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
CHIRONOMIDAE															
Tanypodinae	-	-	-	-	-	-	5	-	-	-	-	-	-	-	-
<u>Cardiocladius</u>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<u>Cricotopus/ Orthocladius</u>	25	-	10	5	5	5	25	5	-	10	-	5	10	-	-
<u>Eukiefferiella/ Tvetenia</u>	-	-	10	-	5	5	15	-	-	-	-	-	-	-	-
<u>Parametrioctenus</u>	-	-	-	-	-	5	-	-	-	-	-	-	-	-	-
<u>Chironomus</u>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<u>Polypedilum aviceps</u>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<u>Polypedilum</u> (all others)	10	10	10	5	5	5	-	-	20	-	-	5	5	5	5
Tanytarsini	10	10	10	10	5	5	10	5	30	-	-	5	10	10	5
TOTAL	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100

## APPENDIX XI: Methods for Calculation of the Nutrient Biotic Index

**Definition:** The Nutrient Biotic Index (Smith, 2005) is a diagnostic measure of stream nutrient enrichment identified by macroinvertebrate taxa. The frequency of occurrences of taxa at varying nutrient concentrations allowed the identification of taxon-specific nutrient optima using a method of weighted averaging. The establishment of nutrient optima is possible based on the observation that most species exhibit unimodal response curves in relation to environmental variables (Jongman et al., 1987). The assignment of tolerance values to taxa based on their nutrient optimum provided the ability to reduce macroinvertebrate community data to a linear scale of eutrophication from oligotrophic to eutrophic. Two tolerance values were assigned to each taxon, one for total phosphorus, and one for nitrate (listed in Smith, 2005). This provides the ability to calculate two different nutrient biotic indices, one for total phosphorus (NBI-P), and one for nitrate (NBI-N). Study of the indices indicate better performance by the NBI-P, with strong correlations to stream nutrient status assessment based on diatom information.

**Calculation of the NBI-P and NBI-N:** Calculation of the indices [2] follows the approach of Hilsenhoff (1987).

$$\text{NBI Score}_{(\text{TP or NO}_3^-)} = \sum (a \times b) / c$$

Where *a* is equal to the number of individuals for each taxon, *b* is the taxon's tolerance value, and *c* is the total number of individuals in the sample for which tolerance values have been assigned.

**Classification of NBI Scores:** NBI scores have been placed on a scale of eutrophication with provisional boundaries between stream trophic status.

Index	Oligotrophic	Mesotrophic	Eutrophic
NBI-P	< 5.0	> 5.0 - 6.0	> 6.0
NBI-N	< 4.5	> 4.5 - 6.0	> 6.0

### References:

- Hilsenhoff, W. L., 1987, An improved biotic index of organic stream pollution. *The Great Lakes Entomologist* 20(1): 31-39.
- Jongman, R. H. G., C. J. F. ter Braak and O. F. R. van Tongeren, 1987, *Data analysis in community and landscape ecology*. Pudoc Wageningen, Netherlands, 299 pages.
- Smith, A.J., 2005, *Development of a Nutrient Biotic Index for use with benthic macroinvertebrates*. SUNY Albany, Masters Thesis, 70 pages.

